



— BUREAU OF —
RECLAMATION

Long-Term Operation

Appendix N – New Melones Stepped Release Plan

Central Valley Project, California

Interior Region 10 – California-Great Basin

Mission Statements

The U.S. Department of the Interior protects and manages the Nation's natural resources and cultural heritage; provides scientific and other information about those resources; honors its trust responsibilities or special commitments to American Indians, Alaska Natives, and affiliated Island Communities.

The mission of the Bureau of Reclamation is to manage, develop, and protect water and related resources in an environmentally and economically sound manner in the interest of the American public.



— BUREAU OF —
RECLAMATION

Long-Term Operation

Appendix N – New Melones Stepped Release Plan

Central Valley Project, California

Interior Region 10 – California-Great Basin

Contents

	Page
1. Introduction.....	1
2. Initial Alternatives Report.....	3
2.1 Management Questions	3
2.2 Initial Analyses.....	3
2.3 Initial Findings	3
2.4 Subsequent Consideration	5
3. Public Draft Environmental Impact Statement Scenarios	7
3.1 Run of River.....	7
3.2 No Action	7
3.3 Alternative 1 – WQCPs.....	7
3.4 Alternative 2 – Multi-Agency Consensus	7
3.5 Alternative 3 – Modified Natural Flow Hydrograph	7
3.6 Alternative 4 – Reservoir Flexibility.....	7
4. Performance Metrics.....	9
4.1 Biological	9
4.2 Water Supply.....	9
4.3 NEPA Resources	10
5. Methods Selection.....	11
5.1 Datasets	11
5.2 Literature	11
5.3 Models.....	27
5.3.1 Water Operations	27
5.3.2 Water Temperature	28
6. Lines of Evidence	29
7. Uncertainty.....	31
8. References.....	33

1. Introduction

The New Melones Stepped Release Plan addresses the volume of instream flows that can occur over a multi-year hydrology without affecting reservoir levels to the extent of depleting the water pool to cause the release of warm water.

This page intentionally left blank

2. Initial Alternatives Report

2.1 Management Questions

The United States Department of the Interior, Bureau of Reclamation's (Reclamation) management questions for the formulation of an alternative include:

- What is the relationship between releases and downstream water temperatures?
- What reservoir storage levels result in the release of warm water?
- Does the long-term instream release result in storage levels that would result in the release of warm water that would affect salmonid survival?
- What risks occur from operating to a 75% exceedance forecast early in the water year?
- What hydrograph shape optimizes Central Valley (CV) steelhead anadromy and survival? Is there a flow intensity threshold to cue migration?
- What is the optimal pulse flow timing by water year type to increase salmonid survival, increase life history diversity, and contribute to successful spawning in the adult population? What migratory phenotypes (i.e., fry, parr, smolts) are more likely to survive under different flow regimes?
- How do releases on the Stanislaus River affect the Water Quality Control Plan (WQCP) and exports?

2.2 Initial Analyses

2.3 Initial Findings

What is the relationship between releases and downstream water temperatures?

- Modeling is under development

What reservoir storage levels result in the release of warm water?

- Modeling is under development

Does the long-term instream release result in storage levels that would result in the release of warm water that would affect salmonid survival?

- Modeling is under development

What risks occur from operating to a 75% exceedance forecast early in the water year?

A sensitivity was run to analyze changes to New Melones operations if different forecast exceedances are used. The SRP uses the San Joaquin River Index to set flows and allocations. The forecasts are only different between February and April, so changes are limited. The 75% forecast is very similar to the final. February, when the forecast is most uncertain is when the forecast is likely to be different than the final water year type under the 75% forecast. Using the 90% forecast, which can reduce flows, results in consistently higher storage (since operations at New Melones are largely independent of storage, the additional storage can stay in the reservoir until evaporation or flood control increases reduce it).

What hydrograph shape optimizes steelhead anadromy and survival? Is there a flow intensity threshold to cue migration?

Spring pulse flows occurring during the steelhead outmigration period are designed to increase life history diversity by cueing anadromy and improving migratory habitat in the Stanislaus River, San Joaquin River, and southern Sacramento–San Joaquin Delta (Delta) (Appendix C, *Species Spatial and Temporal Domains*; NMFS 2009). Spring pulses typically consist of multiple pulses that extend flow variability later into the season, allowing opportunities for broader salmonid outmigration timing. Higher flows cue outmigration and can reduce water temperature and inundate more shallow-water habitat. Maximum pulse flows are generally constrained to 1,500 cfs due to seepage concerns in agricultural lands adjacent to the Stanislaus River (Appendix C).

What is the optimal pulse flow timing by water year type to increase salmonid survival, increase life history diversity, and contribute to successful spawning adult population? What migratory phenotypes (i.e., fry, parr, smolts) are more likely to survive under different flow regimes?

Each year, several pulse flows occur to benefit different life stages of CV steelhead and other salmonids, as specified by Action III.1.3 of the 2009 NMFS Biological Opinion. The 2011 RPA Amendment to Action III.1.3 specifies that pulse flow timing, magnitude, and duration may be adjusted based on water year type, with longer durations of pulses and higher intensities of peaks during wetter years, and other adjustments based on a variety of other considerations (Table 4). Maximum pulse flows are generally limited to 1,500 cfs and a duration of 10 days due to seepage concerns in agricultural lands adjacent to the Stanislaus River (Table 4).

Fall pulse flows usually occur in October through early November prior to peak spawning by adults, and timing may vary based on environmental conditions to achieve suitable habitat (DO and temperature) conditions for migration and holding. These multi-peak (3-4 peaks) flow pulses are designed to attract adult CV steelhead to the Stanislaus while deterring adults from spawning in areas that may be subsequently de-watered (NMFS 2009; Appendix A, *Facilities Description*).

Winter instability flows usually occur between January and February (may occur in both months) and may be timed to coincide with winter storms, as well as with emergence of fall-run Chinook fry. These flows are shaped to simulate winter storms, with rapidly increasing flows followed by slowly descending flows. Higher flows may inundate shallow water habitat, leading to benefits for rearing juvenile salmonids that may take refuge from predators and receive potential growth benefits from increased terrestrial inputs (Appendix B, *Water Operations and Ecosystem Analyses*).

Spring pulse flows usually occur between April and June. These multi-peak (3-6 peaks) flow pulses are designed to increase flow variability to allow for a broader range of salmonid outmigration timing, increasing life history diversity. These flows are intended to cue anadromy of steelhead and outmigration of smolts and improve migratory habitat downstream, as well as convey outmigrating salmonids (smolts) more rapidly. Pulse flows may also inundate shallow water habitat, leading to benefits for rearing salmonids that may take refuge from predators and receive potential growth benefits (Appendix C).

Sturrock et al. (2020) found that greater flow volume and variability was associated with earlier outmigration, and thus a greater proportion of fish outmigrating as fry. Winter flows had a large impact on whether or not fry outmigrated downstream, with reduced winter flows leading to later salmon outmigration, composed of smaller numbers of larger fish, and ultimately leading to lower production. Annual abundances of emigrant per spawner were correlated with flow variability and mean rearing flows across all migratory phenotypes. In general, while fry and smolts were the more dominant phenotypes for outmigration, survival was higher for intermediate-sized (parr) fish outmigrating in the spring.

How do releases on the Stanislaus River affect WQCP and exports?

There are only small impacts to the WQCP and exports from the Stanislaus River Initial Alternatives.

2.4 Subsequent Consideration

This page intentionally left blank

3. Public Draft Environmental Impact Statement Scenarios

Under the National Environmental Policy Act, Reclamation compares action alternatives to a “no action” alternative. Under the Endangered Species Act, Reclamation’s discretionary actions over an environmental baseline determine the effects on listed species. No single environmental baseline to evaluate the effects under the Endangered Species Act (ESA) or impacts under NEPA. ESA requires a comparison to the environmental baseline which is informed by ROR and Alt 1. NEPA requires a comparison to NA.

3.1 Run of River

[Placeholder]

3.2 No Action

[Placeholder]

3.3 Alternative 1 – WQCPs

[Placeholder]

3.4 Alternative 2 – Multi-Agency Consensus

[Placeholder]

3.5 Alternative 3 – Modified Natural Flow Hydrograph

[Placeholder]

3.6 Alternative 4 – Reservoir Flexibility

[Placeholder]

This page intentionally left blank

4. Performance Metrics

Performance metrics describe criteria that can be measured, estimated, or calculated relevant to informing trade-offs for alternative management actions.

4.1 Biological

Fisheries metrics consider direct observations and environmental surrogates including:

- **Water Temperature for Juveniles and Eggs**
 - The survival temperature threshold for juvenile steelhead is less than or equal to 68 degrees Fahrenheit (°F) from May 1 to October 31
 - The temperature threshold for steelhead egg incubation is less than or equal to 54°F from December 1 to May 31

The steelhead on the Stanislaus River generally move upstream to spawn between July and March; juvenile steelhead out-migrate between January and June (NMFS 2014). Reservoir releases combined with other environmental conditions will affect water temperature, dissolved oxygen level, and other habitat attributes that will influence the timing, condition and survival of eggs in the spawning redds. Decrease in flows generally results in warmer water temperatures in the winter; however, winter water temperatures are colder than adult migration temperature needs. The New Melones Stepped Release Plan promotes increased storage at New Melones Reservoir, which can help the development of a larger cold water pool. More cold water in the reservoir may help lower water temperatures downstream of Goodwin Dam, which would benefit steelhead in all life stages in the lower Stanislaus River.

The fisheries metrics address the volume of instream flows that can occur over a multi-year hydrology without affecting reservoir levels to the extent of depleting the cold water pool to cause the release of warm water.

4.2 Water Supply

Water supply metrics consider the multipurpose beneficial uses of New Melones including:

- Central Valley Project (CVP) water service contracts
- Senior water right holders (Oakdale Irrigation District and South San Joaquin Irrigation District)
- State Water Resources Control Board Water Right Decision 1641 (D-1641) standards and its dissolved oxygen requirement
- Flood conservation pool release

CalSim II would support the evaluation of water supply metrics.

4.3 NEPA Resources

Analysis of the range of alternatives as required by the National Environmental Policy Act is anticipated to describe changes in the multiple resources areas. Key resources are anticipated to include surface water supply, water quality, groundwater resources, aquatic resources, terrestrial biological resources, regional economics, land use and agricultural resources, cultural resources, socioeconomics, environmental justice, climate change, and power generation.

5. Methods Selection

Reclamation solicited input from the stakeholders and agencies for the knowledge base paper (Steelhead Juvenile Production Estimate). Reclamation identified the following datasets, literature, and models to help in evaluating the New Melones Stepped Release Plan.

5.1 Datasets

Several efforts to characterize historical and ongoing steelhead monitoring programs in the California Central Valley have been completed over the past two decades. A few years after the completion of the Central Valley Steelhead Monitoring Plan, a series of related monitoring projects, identified as the Central Valley Steelhead Monitoring Program, were initiated on the Sacramento River and its tributaries (Fortier et al. 2014). Appendix G, *Specific Facility and Water Operations Deconstruction*, provides pertinent information for the datasets for steelhead.

CalFish (2019). CalFish – A California cooperative anadromous fish and habitat data program. Middle Sacramento River salmon and Steelhead monitoring. Available: <https://www.calfish.org/ProgramsData/ConservationandManagement/CentralValleyMonitoring/SacramentoValleyTributaryMonitoring/MiddleSacramentoRiverSalmonandSteelheadMonitoring.aspx>.

SacPAS: Central Valley Prediction & Assessment of Salmon provides a platform for data queries of juvenile steelhead salvage and loss. Available: http://www.cbr.washington.edu/sacramento/data/juv_salvage_loss.html.

Use CalFishTrack to understand juvenile steelhead routing and survival into the Delta. <https://oceanview.pfeg.noaa.gov/CalFishTrack/>

5.2 Literature

The documents listed below were compiled from the 2019 BiOps, 2020 ITP, fact sheets produced for the February 2021 joint Delta Science Program – U.S. Bureau of Reclamation Steelhead Workshop, and a Google Scholar search.

Abadía-Cardoso, A., A. J. Clemento, and J. C. Garza. 2011. Discovery and characterization of single- nucleotide polymorphisms in steelhead/rainbow trout, *Oncorhynchus mykiss*. *Molecular Ecology Resources* 11:31–49.

Adams, B. L., W. S. Zaugg, and L. R. McLain. 1975. Inhibition of Salt Water Survival and Na-KATPase Elevation in Steelhead Trout (*Salmo gairdneri*) by Moderate Water Temperatures. *Transactions of the American Fisheries Society* 104(4):766–769.

- Alston, N. O., J. M. Newton, and M. R. Brown. 2007. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from November 2003 through November 2004. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Beakes, M. P., W. H. Satterthwaite, E. M. Collins, D. R. Swank, J. E. Merz, R.G. Titus, S.M. Sogard, and M. Mangel, 2010. Smolt transformation in two California steelhead populations: effects of temporal variability in growth. *Transactions of the American Fisheries Society* 139(5) 1263–1275.
- Beakes, M. P., J. W. Moore, S. A. Hayes, and S. M. Sogard. 2014. Wildfire and the effects of shifting stream temperature on salmonids. *Ecosphere* 5(5):1–14.
- Beakes, M. P., S. Sharron, R. Charish, J. W. Moore, W. H. Satterthwaite, E. Sturm, B. K. Wells, S. M. Sogard, and M. Mange. 2014. Using scale characteristics and water temperature to reconstruct growth rates of juvenile steelhead *Oncorhynchus mykiss*. *Journal of Fish Biology* 84(1):58–72.
- Beakes, M., R. Bilski, B. Mattias, B. Byrne, P. Vick, and P. Goertler. 2021. *Oncorhynchus mykiss* Monitoring and Research Gap Analysis. In Monitoring Steelhead Populations in the San Joaquin Basin. Edited by Delta Science Program - Delta Stewardship Council, Joint Delta Science Program – U.S. Bureau of Reclamation Workshop, Sacramento, CA. DOI: 10.13140/RG.2.2.29383.21927 (<https://deltacouncil.ca.gov/pdf/science-program/factsheets/2021-02-03-monitoring-steelhead-populations-monitoring-and-research-gap-analysis.pdf>)
- Beakes, M., and C. Phillis. 2021. Life-History Variation in *Oncorhynchus mykiss*. In Monitoring Steelhead Populations in the San Joaquin Basin. Edited by Delta Science Program - Delta Stewardship Council, Joint Delta Science Program – U.S. Bureau of Reclamation Workshop, Sacramento, CA. DOI: 10.13140/RG.2.2.15438.79685 (<https://deltacouncil.ca.gov/pdf/science-program/fact-sheets/2021-02-04-monitoringsteelhead-populations-life-history-variation.pdf>).
- Berejikian, B. A., L. A. Campbell, and M. E. Moore. 2013. Large-scale freshwater habitat features influence the degree of anadromy in eight Hood Canal *Oncorhynchus mykiss* populations. *Canadian Journal of Fisheries and Aquatic Sciences* 70(5):756–765.
- Bjornn, T. C. 1978. Survival, production, and yield of trout and chinook salmon in the Lemhi River, Idaho A Final Report for Federal Aid to Fish Restoration, Project F-49-R, Salmon and Steelhead Investigations. Idaho Cooperative Fishery Research Unit.
- Boughton, D. A., and A. S. Pike. 2013. Floodplain rehabilitation as a hedge against hydroclimatic uncertainty in a migration corridor of threatened steelhead. *Conservation Biology* 27(6):1158–1168.

- Bottaro, R. J., L. A. Earley, and M. R. Brown. 2012. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2010. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and M. R. Brown. 2012. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2011. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., L. A. Earley, and M. R. Brown. 2013. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2012. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., L. A. Earley, and M. R. Brown. 2014. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2013. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and M. R. Brown. 2018. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2014. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and M. R. Brown. 2018. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2015. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and L. A. Earley. 2018. Monitoring Adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2016. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and L. A. Earley. 2019. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2017. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Bottaro, R. J., and L. A. Earley. 2020. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2018. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Brown, M. R. 1996. Benefits of Increased Minimum Instream Flows on Chinook Salmon and Steelhead in Clear Creek, Shasta County, California 1995-6, USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Brown, M. R., and J. M. Newton. 2002. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through October 2001. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.

- Brown, M. R., N. O. Alston, and J. M. Newton. 2005. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2002. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Brown, M. R., and N. O. Alston. 2007. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from November 2002 through November 2003. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Buchanan, R. A., E. Buttermore, and J. Israel. 2021. Outmigration survival of a threatened steelhead population through a tidal estuary. *Canadian Journal of Fisheries and Aquatic Sciences* 78(12):1869–1886.
- Cada, G. F., M. D. Deacon, S. V. Mitz, and M. S. Bevelhimer. 1997. Effects of water velocity on the survival of downstream-migrating juvenile salmon and steelhead: A review with emphasis on the Columbia River basin. *Reviews in Fisheries Science* 5(2):131–183.
- California Department of Water Resources. 2008. Quantification of Pre-Screen Loss of Juvenile Steelhead within Clifton Court Forebay. Page 136. Fishery Improvements Section, editor.
- California Department of Fish and Wildlife. 2016. Memorandum to Amanda Cranford, National Marine Fisheries Service from Rob Titus, California Department of Fish and Wildlife, Trends in Central Valley Steelhead Harvest. Page 3. California Department of Fish and Wildlife, editor.
- California Department of Water Resources. 2018. Effects of the South Delta Agricultural Barriers on Emigrating Juvenile Salmonids. Page 244. California Department of Water Resources, editor, Sacramento, CA.
- Carlson, S. M., and T. R. Seamons. 2008. A review of quantitative genetic components of fitness in salmonids: implications for adaptation to future change. *Evolutionary Applications* 1(2):222–238.
- Carter, K. 2005. The Effects of Dissolved Oxygen on Steelhead Trout, Coho Salmon, and Chinook Salmon Biology and Function by Life Stage. Page 10. California Regional Water Quality Control Board, editor, North Coast Region.
- Cech Jr, J., and C. A. Myrick. 1999. Steelhead and Chinook Salmon Bioenergetics: Temperature, Ration, and Genetic Effects. University of California, Davis.
- Chapman, E. D., Hearn, A.R., Singer, G.P., Brostoff, W.N., LaCivita, P.E. and A.P. Klimley. 2015. Movements of steelhead (*Oncorhynchus mykiss*) smolts migrating through the San Francisco Bay Estuary. *Environmental Biology of Fishes* 98(4):1069–1080.

- Chapman, E. P., Hearn, A.R., Michel, C.J., Ammann, A.J., Lindley, S.T., Thomas, M.J., Sandstrom, P.T., Singer, G.P., Peterson, M.L., MacFarlane, R.B. and A.P. Klimley. 2013. Diel movements of Out-migrating Chinook salmon (*Oncorhynchus tshawytscha*) and steelhead trout (*Oncorhynchus mykiss*) smolts in the Sacramento/San Joaquin watershed. *Environmental Biology of Fishes* 96:273–286.
- Clark, K. W., M. D. Bowen, R. B. Mayfield, K. P. Zehfuss, J. D. Taplin, and C. H. Hanson. 2009. Quantification of pre-screen loss of juvenile steelhead in Clifton Court Forebay. California Department of Water Resources, Bay-Delta Office, Fishery Improvements Section, Sacramento, CA. March 2009.
- Colby, D. J., and M. R. Brown. 2012. Juvenile salmonid monitoring in Battle Creek, California, November 2010 through June 2011. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Courter, I., C. Justice, and S. Cramer. 2009. Flow and temperature effects on life history diversity of *Oncorhynchus mykiss* in the Yakima River basin. Cramer Fish Sciences, Gresham, Oregon.
- Courter, I. I., Courter, L., Garrison, T., Cramer, D., Duery, S., Child, D., Hanna, T. and E. Buckner. 2012. Effects of the Aquatic Herbicide Cascade on survival of Salmon and Steelhead Smolts during Seawater transition.
- Earley, J. T., D. J. Colby, and M. R. Brown. 2008. Juvenile salmonid monitoring in Clear Creek, California, from July 2002 through September 2003. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, California
- Earley, J. T., D. J. Colby, and M. R. Brown. 2008. Juvenile salmonid monitoring in Clear Creek, California, from July 2002 through September 2003. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Earley J.T., D.J. Colby, and M.R. Brown. 2008. Juvenile Salmonid Monitoring in Clear Creek, California, from October 2003 through September 2004. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Earley, J. T., D. J. Colby, and M.R. Brown. 2008. Juvenile salmonid monitoring in Clear Creek, California, from October 2006 through September 2007. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Earley, J. T., D. J. Colby, and M. R. Brown. 2009. Juvenile salmonid monitoring in Clear Creek, California, from October 2007 through September 2008. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Earley, J. T., D. J. Colby, and M. R. Brown. 2010. Juvenile salmonid monitoring in Clear Creek, California, from October 2008 through September 2009. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.

- Earley, J. T., D. J. Colby, and M. R. Brown. 2013. Juvenile salmonid monitoring in Clear Creek, California, from October 2009 through September 2010. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Earley, J. T., D. J. Colby, and M. R. Brown. 2013. Juvenile salmonid monitoring in Clear Creek, California, from October 2010 through September 2011. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Ebersole, J. L., W. J. Liss, and C. A. Frissell. 2001. Relationship between stream temperature, thermal refugia and rainbow trout *Oncorhynchus mykiss* abundance in arid-land streams in the northwestern United States. *Ecology of Freshwater Fish* 10:1–10.
- Eilers, C. D. 2010. Review of present steelhead monitoring programs in the California Central Valley May 2008. Fisheries Branch Administrative Report Number: 2010-1
- Ellrott, B., H. Brown, and R. Johnson. California Central Valley Steelhead Distinct Population Segment. In *Monitoring Steelhead Populations in the San Joaquin Basin*. Edited by Delta Science Program - Delta Stewardship Council, Joint Delta Science Program – U.S. Bureau of Reclamation Workshop, Sacramento, CA. DOI: 10.13140/RG.2.2.30997.04329 (<https://deltacouncil.ca.gov/pdf/science-program/fact-sheets/2021-02-04-california-central-valley-steelhead-distinct-population-segment-factsheet.pdf>)
- Eschenroeder, J. C., M. L. Peterson, M. Hellmair, T. J. Pilger, D. Demko, and A. Fuller. 2022. Counting the Parts to Understand the Whole: Rethinking Monitoring of Steelhead in California’s Central Valley. *San Francisco Estuary and Watershed Science* 20(1).
- Fortier, R., J. Nelson, R. Bellmer, and R. Nielson. 2014. Implementation Plan for the Central Valley Steelhead Monitoring Program. California Department of Fish and Wildlife, Fisheries Branch. July 2014.
- Gaines, P. D., and C.D. Martin. 2001. Abundance and seasonal, spatial and diel distribution patterns of juvenile salmonids passing the Red Bluff Diversion Dam, Sacramento River. Red Bluff Research Pumping Plant Report Series, Volume 14. U.S. Fish and Wildlife Service, Red Bluff, CA.
- Gaines, P. D., R. E. Null and M. R. Brown. 2003. Estimating the abundance of Clear Creek juvenile chinook salmon and steelhead trout by use of a rotary-screw trap. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office. Progress Report (Vol.1), February 2003.
- Giovannetti, S. L., M. R. Brown. 2007. Central Valley Steelhead and Late Fall Chinook Salmon Redd Surveys on Clear Creek, California 2007. U.S. Fish and Wildlife Service, Red Bluff, CA.
- Giovannetti, S. L., M. R. Brown. 2008. Steelhead and Late-fall Chinook Salmon Redd Surveys on Clear Creek, California 2008 Annual Report. U.S. Fish and Wildlife Service, Red Bluff, CA.

- Giovannetti, S.L., M. R. Brown. 2009. Adult Steelhead and Late-fall Chinook Salmon Monitoring on Clear Creek, California. 2009 Annual Report. U.S. Fish and Wildlife Service, Red Bluff, CA.
- Giovannetti, S. L., R. J. Bottaro, and M. R. Brown. 2013. Adult steelhead and late-fall Chinook salmon Monitoring on Clear Creek, California: 2010 Annual report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Giovannetti, S. L., R. J. Bottaro, and M. R. Brown. 2013. Adult steelhead and late-fall Chinook salmon Monitoring on Clear Creek, California: 2011 Annual report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Good, T. P., R. S. Waples, and P. Adams. 2005. Updated status of federally listed ESUs of West Coast salmon and steelhead. National Oceanic and Atmospheric Administration, National Marine Fisheries Service. NOAA Technical Memorandum NMFS-NWFSC-66. June 2005.
- Greenwald, G. M., J. T. Earley, and M. R. Brown. 2003. Juvenile salmonid monitoring in Clear Creek, California, from July 2001 to July 2002. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Hallock, R. J., D. H. Fry Jr., and D. A. LaFauce. 1957. The Use of Wire Fyke Traps to Estimate the Runs of Adult Salmon and Steelhead in the Sacramento River. *California Fish and Game* 43(4):271-298.
- Hallock, R. J., W. F. Van Woert, and L. Shapovalov. 1961. An Evaluation of Stocking Hatchery-reared Steelhead Rainbow Trout (*Salmo gairdnerii gairdnerii*) in the Sacramento River System. *Fish Bulletin* 114:3-74.
- Hannon, J. 2013. American River Steelhead (*Oncorhynchus mykiss*) Spawning - 2013, with Comparison to Prior Years. Pages 1-32. U.S. Bureau of Reclamation.
- Hannon, J., and B. Deason. 2008. American River Steelhead (*Oncorhynchus mykiss*) Spawning 2001-2007. U.S. Department of the Interior, Bureau of Reclamation, Mid-Pacific Region.
- Hannon, J., M. Healey, and B. Deason. 2003. American River Steelhead (*Oncorhynchus mykiss*) Spawning 2001-2003. Pages 36. U.S. Bureau of Reclamation, and California Department of Fish and Game, Sacramento, California.
- Hayes, S. A., M. H. Bond, C. V. Hanson, A.W. Jones, A. J. Ammann, J. A. Harding, A. L. Collins, J. Perez, and R. B. MacFarlane. 2011. Down, up, down and “smolting” twice? Seasonal movement patterns by juvenile steelhead (*Oncorhynchus mykiss*) in a coastal watershed with a bar closing estuary. *Canadian Journal of Fisheries and Aquatic Sciences* 68(8):1341-1350.
- Hodge, B. W., M. A. Wilzbach, W. G. Duffy, R. M. Quiñones, and J. A. Hobbs. 2016. Life history diversity in Klamath River steelhead. *Transactions of the American Fisheries Society* 145(2): 227-238.

- Hutchings, J. A. 2011. Old wine in new bottles: reaction norms in salmonid fishes. *Heredity* 106(3):421–437.
- IEP. 1999. Monitoring, Assessment, and Research on Central Valley Steelhead: Status of Knowledge, Review of Existing Programs, and Assessment of Needs. Interagency Ecological Program Steelhead Project Work Team.
- Israel, J., H. Brown, and L. Conrad. 2021. Workshop Objectives and Regulatory Background. In Monitoring Steelhead Populations in the San Joaquin Basin. Edited by Delta Science Program - Stewardship Council, Joint Delta Science Program – U.S. Bureau of Reclamation Workshop, Sacramento, CA. DOI: 10.13140/RG.2.2.33390.33606 (<https://deltacouncil.ca.gov/pdf/science-program/fact-sheets/2021-02-03-monitoring-steelhead-populations-workshop-objectives-and-regulatory-background.pdf>)
- Karp, C., B. Wu, and K. Kumagai. 2017. Juvenile Chinook Salmon, Steelhead, and Adult Striped Bass Movements and Facility Efficiency at the Tracy Fish Collection Facility Tracy Technical Bulletin 2017-1. Pages 81
- Keefer, M. L., C. A. Peery, and B. High. 2009. Behavioral thermoregulation and associated mortality trade-offs in migrating adult steelhead (*Oncorhynchus mykiss*): variability among sympatric populations. *Canadian Journal Fisheries Aquatic Science* 66:1734–1747.
- Keeley, E. R. 2001. Demographic responses to food and space competition. *Ecology* 82:1247–1259.
- Kelson, S. J., M. R. Miller, T. Q. Thompson, S. M. O'Rourke, and S. M. Carlson. 2019. Do genomics and sex predict migration in a partially migratory salmonid fish, *Oncorhynchus mykiss*? *Canadian Journal of Fisheries and Aquatic Sciences* 76(11):2080–2088.
- Kelson, S.J., S. M. Carlson, and M. R. Miller. 2020. Indirect genetic control of migration in a salmonid fish. *Biology Letters* 16(8), p.20200299.
- Kendall, N.W., J. R. McMillan, M. R. Sloat, T. W. Buehrens, T. P. Quinn, G. R. Pess, K.V Kuzishchin, M. M. McClure, and R. W. Zabel. 2015. Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): a review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* 72(3):319–342.
- Lee, D. P., and J. Chilton. 2007. Hatchery and Genetic Management Plan for Nimbus Fish Hatchery Winter-Run Steelhead Program.
- Lindley, S.T., R. Schick, E. Mora, P.B. Adams, J.J. Anderson, S. Greene, C. Hanson, B.P. May, D.R. McEwan, R.B. MacFarlane, C. Swanson, and J.G. Williams (2007). Framework for assessing viability of threatened and endangered Chinook salmon and steelhead in the Sacramento-San Joaquin Basin. *San Francisco Estuary and Watershed Science* 5(1).

- Louie, S., R. Henery, J. Rosenfield, J. Shelton, J. Zimmerman, M. Gutierrez, B. Ellrott, J. Wikert, J. Howard, T. Heyne, and A. Weber-Stover. 2019. Conservation Planning Foundation for Restoring Chinook Salmon (*Oncorhynchus Tshawytscha*) and *O. mykiss* in the Stanislaus River.
- Lufkin, A., editor. 1991. *California's Salmon and Steelhead: The Struggle to Restore an Imperiled Resource*. Berkeley: University of California Press, Berkeley.
- Mangel, M., and W. H. Satterthwaite. 2008. Combining proximate and ultimate approaches to understand life history variation in salmonids with application to fisheries, conservation, and aquaculture. *Bulletin of Marine Science* 83(1):107–130.
- McEwan, D., and T. A. Jackson. 1996. Steelhead Restoration and Management Plan for California. Pages 1-234.
- McMillan, J., S. L. Katz, and G. R. Pess. 2007. Observational evidence of spatial and temporal structure in a sympatric anadromous (winter steelhead) and resident rainbow trout mating system on the Olympic Peninsula, Washington. *Transactions of the American Fisheries Society* 136(3):736–748.
- McMillan, J. R., J. Dunham, G. H. Reeves, J. S. Mills, and C. E. Jordan. 2012. Individual condition and stream temperature influence early maturation of rainbow and steelhead trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes* 93:343–355. doi:10.1007/s10641-011-9921-0.
- Miller, M. R., J. P. Brunelli, P. A. Wheeler, S. Liu, C. E. Rexroad III, Y. Palti, C. Q. Doe, and G. H. Thorgaard. 2012. A conserved haplotype controls parallel adaptation in geographically distant salmonid populations. *Molecular Ecology* 21(2):237–249.
- Moyle, P. B. 2002. *Inland Fishes of California*, 2nd edition. Berkeley, CA: University of California Press.
- Moffitt, 2014. Histological assessment of organs in sexually mature and post-spawning steelhead trout and insights into iteroparity. *Rev Fish Biol Fisheries* (2014) 24:781–801.
- Myrick, C., and J. Cech Jr. 2000. Growth and thermal biology of Feather River steelhead under constant and cyclical temperatures. University of California, Davis, Department of Wildlife, Fish, and Conservation Biology.
- Myrick, C. A., and J. J. Cech. 2001. Temperature effects on Chinook salmon and steelhead: a review focusing on California's Central Valley populations. Bay-Delta Modeling Forum.
- Myrick, C. A., and J. J. Cech Jr. 2005. Effects of temperature on the growth, food consumption, and thermal tolerance of age-0 Nimbus-strain steelhead. *North American Journal of Aquaculture* 67:324–330.

- Narum S. R., D. Hatch, A. J. Talbot, P. Moran, and M. Powell, 2008. Iteroparity in complex mating systems of steelhead *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Biology* 72(1):45–60
- National Marine Fisheries Service. 2011. 5-Year Review: Summary and Evaluation of Central Valley Steelhead Distinct Population Segment.
- National Marine Fisheries Service. 2014. Final recovery plan for the evolutionarily significant units of Sacramento River winter-run Chinook salmon and Central Valley spring-run Chinook salmon and the distinct population segment of California Central Valley steelhead. National Oceanic and Atmospheric Administration, National Marine Fisheries Service. July 2014.
- National Marine Fisheries Service. 2016a. 5-Year Status Review: Summary and Evaluation of California Central Valley Steelhead Distinct Population Segment. Sacramento, California.
- National Marine Fisheries Service. 2016b. 2016 5-Year Review: Summary & Evaluation of Central California Coast Steelhead. West Coast Region.
- Newton, J. M., and M. R. Brown. 2003. Middle Battle Creek reconnaissance survey, 2001. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Newton, J. M., N. O. Alston, and M. R. Brown. 2007. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from November 2004 through November 2005. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Newton, J. M., N. O. Alston, and M. R. Brown. 2007. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2006. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Newton, J. M., L. A. Stafford, and M. R. Brown. 2008. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2007. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Newton, J. M., and L.A. Stafford. 2011. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2009. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Nichols, K. M., A. F. Edo, P. A. Wheeler, and G. H. Thorgaard. 2008. The genetic basis of smoltification-related traits in *Oncorhynchus mykiss*. *Genetics* 179(3):1559–1575.
- Nielsen, J. L., T. E. Lisle, and V. Ozaki. 1994. Thermally Stratified Pools and Their Use by Steelhead in Northern California Streams. *Transactions of the American Fisheries Society* 123(4):613–626.

- Nielson, J. L., S. Pavey, T. Wiacek, G. K. Sage, and I. Williams. 2003. Genetic Analyses of Central Valley Trout Populations 1999–2003. California Department of Fish and Game and U. S. Fish and Wildlife Service, 44 pp.
- Nobriga, M. L., and P. Cadrett. 2001. Differences among Hatchery and Wild Steelhead; Evidence from Delta Fish Monitoring Programs Pages 56 in IEP Newsletter Summer.
- Ohms, H. A., M. R. Sloat, G. H. Reeves, C. E. Jordan, and J. B. Dunham. 2014. Influence of sex, migration distance, and latitude on life history expression in steelhead and rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences* 71(1):70–80.
- Pearse, D. E., M. R. Miller, A. Abadía-Cardoso, and J. C. Garza. 2014. Rapid parallel evolution of standing variation in a single, complex, genomic region is associated with life history in steelhead/rainbow trout. *Proceedings of the Royal Society B: Biological Sciences*, 281(1783), p.20140012.
- Pearse, D. E., and J. C. Garza. 2015. You can't unscramble an egg: Population genetic structure of *Oncorhynchus mykiss* in the California Central Valley inferred from combined microsatellite and single nucleotide polymorphism data. *San Francisco Estuary and Watershed Science* 13(4).
- Pearsons, T. N., G. M. Temple, A. L. Fritts, C. L. Johnson, and T. D. Webster. 2008. Ecological interactions between non-target taxa of concern and hatchery supplemented salmon. 2007 Annual Report Project, (1995-063), p.25.
- Phillips, R. W., and H. J. Campbell. 1961. The Embryonic Survival of Coho Salmon and Steelhead Trout as Influenced by Some Environmental Conditions in Gravel Beds. Pages 60-72 in Fourteenth annual report. Pacific Marine Fisheries Commission, Portland, Oregon.
- Phillis, C. C., J. W. Moore, M. Buoro, S. A. Hayes, J. C. Garza, and D. E. Pearse. 2016. Shifting thresholds: rapid evolution of migratory life histories in steelhead/rainbow trout, *Oncorhynchus mykiss*. *Journal of Heredity* 107(1):51–60.
- Poytress, W. R., J. J. Gruber, F. D. Carrillo, and S. D. Voss. 2014. Compendium Report of Red Bluff Diversion Dam Rotary Trap Juvenile Anadromous Fish Production Indices for Years 2002-2012. Report of U.S. Fish and Wildlife Service to California Department of Fish and Wildlife and U.S. Bureau of Reclamation.
- Provins, S. S. and C. D. Chamberlain 2019. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek, winter 2011 to spring 2012. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Provins S.S. and C.D. Chamberlain 2019. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek; winter 2012 to spring 2013. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.

- Provins S.S. and C.D. Chamberlain. 2019. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek; winter 2013 to spring 2014. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Provins, S.S. and C.D. Chamberlain 2019. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek; winter 2014 to spring 2015. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Provins S.S. and C.D. Chamberlain. 2019. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek; winter 2016 to spring 2017. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Provins S. S. and C. D. Chamberlain. 2020. Distribution and abundance of Rainbow Trout/steelhead and late-fall run Chinook Salmon redds in Clear Creek; winter 2017 to spring 2018. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Raquel, P. F. 1989. Effects of Handling and Trucking on Chinook salmon, striped bass, American Shad, Steelhead Trout, Threadfin Shad, and White Catfish salvaged at the John E. Skinner Delta Fish Protective Facility.
- Richter, A., and S. A. Kolmes. 2005. Maximum Temperature Limits for Chinook, Coho, and Chum Salmon, and Steelhead Trout in the Pacific Northwest. *Reviews in Fisheries Science* 13(1):23–49.
- Rombough, P. J. 1988. Growth, Aerobic Metabolism, and Dissolved-Oxygen Requirements of Embryos and Alevins of Steelhead, *Salmo-Gairdneri*. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 66(3):651–660.
- Sasaki, S. 1966. Distribution and Food Habits of king salmon (*Oncorhynchus tshawytscha*) and steelhead rainbow trout, *Salmo gairdnerii*, in the Sacramento-San Joaquin Delta. *Fish Bulletin* 136:108–114.
- Satterthwaite, W.H., M.P. Beakes, E. M. Collins, D. R. Swank, J. E. Merz, R. G. Titus, S. M. Sogard, and M. Mangel. 2009. Steelhead life history on California's central coast: insights from a state-dependent model. *Transactions of the American Fisheries Society* 138(3):532–548.
- Satterthwaite, W.H., M. P. Beakes, E. M. Collins, D. R. Swank, J. E. Merz, R. G. Titus, S. M. Sogard, and M. Mangel. 2010. State-dependent life history models in a changing (and regulated) environment: steelhead in the California Central Valley. *Evolutionary Applications* 3(3):221–243.
- Schaefer, R. A., S. L. Gallagher, and C. D. Chamberlain. 2019. Distribution and Abundance of California Central Valley steelhead/Rainbow Trout and Late-fall Chinook Salmon Redds in Clear Creek, Winter 2015 to Spring 2016. Red Bluff, CA.

- Schaefer, R. A., S. L. Gallagher, and C.D. Chamberlain. 2019. Distribution and abundance of California Central Valley steelhead/Rainbow Trout and late-fall Chinook Salmon redds in Clear Creek, winter 2015 to spring 2016. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml C. M., J. T. Earley, and L. A. Earley. 2018. Brood Year 2011 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., J. T. Earley, and L. A. Earley. 2019. Brood Year 2012 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and L. A. Earley. 2019. Brood Year 2013 Juvenile Salmonid Monitoring in Battle, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., and L. A. Earley. 2019. Brood Year 2014 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and L. A. Earley. 2019. Brood Year 2015 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and L. A. Earley. 2019. Brood Year 2016 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and L. A. Earley. 2020. Brood Year 2016 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and L. A. Earley. 2021. Brood Year 2018 Juvenile Salmonid Monitoring in Battle Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., J. T. Earley, and C. D. Chamberlain. 2018. Brood Year 2011 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., and C. D. Chamberlain. 2019. Brood Year 2012 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., and C. D. Chamberlain. 2019. Brood Year 2013 Juvenile Salmonid Monitoring in Clear Creek. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.

- Schraml, C. M. and C. D. Chamberlain. 2019. Brood Year 2014 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and C. D. Chamberlain. 2019. Brood Year 2015 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M. and C. D. Chamberlain. 2020. Brood Year 2016 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., J. R. Knight, and C. D. Chamberlain. 2020. Brood Year 2017 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Schraml, C. M., and C. D. Chamberlain. 2021. Brood Year 2018 Juvenile Salmonid Monitoring in Clear Creek, California. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Shapovalov, L., and A. C. Taft. 1954. The life histories of the steelhead rainbow trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus kisutch*): with special reference to Waddell Creek, California, and recommendations regarding their management (p. 575). Sacramento, California, USA: California Department of Fish and Game.
- Shelton, A. O., W. H. Satterthwaite, M. P. Beakes, S. B. Munch, S. M. Sogard, and M. Mangel. 2013. Separating intrinsic and environmental contributions to growth and their population consequences. *The American Naturalist* 181(6):799–814.
- Sigler, J. W., T. C. Bjornn, and F. H. Everest. 1984. Effects of Chronic Turbidity on Density and Growth of Steelheads and Coho Salmon. *Transactions of the American Fisheries Society* 113(2):142–150.
- Sloat, M. R., and A. M. K. Osterback. 2013. Maximum stream temperature and the occurrence, abundance, and behavior of steelhead trout (*Oncorhynchus mykiss*) in a southern California stream. *Canadian Journal of Fisheries and Aquatic Sciences* 70(1):64–73.
- Sloat, M. R., and G. H. Reeves. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* 71(4):491–501.
- Sogard, S. M., J. E. Merz, W. H. Satterthwaite, M. P. Beakes, D. R. Swank, E. M. Collins, R. G. Titus, and M. Mangel. 2012. Contrasts in habitat characteristics and life history patterns of *Oncorhynchus mykiss* in California's Central Coast and Central Valley. *Transactions of the American Fisheries Society* 141(3):747–760.

- Stafford, L.A., and J.M. Newton. 2010. Monitoring adult Chinook salmon, rainbow trout, and steelhead in Battle Creek, California, from March through November 2008. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Stanley, C. E., R. J. Bottaro, and L. A. Earley. 2020. Monitoring adult Chinook Salmon, Rainbow Trout, and Steelhead in Battle Creek, California, from March through November 2019. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Teo, S.L. and multiple co-authors, 2013. Archival and acoustic tags reveal the post-spawning migrations, diving behavior, and thermal habitat of hatchery-origin Sacramento River steelhead kelts. *Environmental Biology of Fishes* 96. DOI:10.1007/s10641-011-9938-4.
- Thrower, F. P., J. J. Hard, and J. E. Joyce. 2004. Genetic architecture of growth and early life-history transitions in anadromous and derived freshwater populations of steelhead. *Journal of Fish Biology* 65:286–307.
- Turner, M. A., M. R. Viant, S. J. Teh, and M. L. Johnson. 2007. Developmental rates, structural asymmetry, and metabolic fingerprints of steelhead trout (*Oncorhynchus mykiss*) eggs incubated at two temperatures. *Fish Physiology and Biochemistry* 33(1):59–72.
- U.S. Bureau of Reclamation (USBR). 2018. NMFS Biological Opinion RPA IV.2.2: 201.1 Six-Year Acoustic Telemetry Steelhead Study. Contributions by Buchanan, R., J. Israel, P. Brandes. E. Buttermore. Reclamation Bay-Delta Office, Mid-Pacific Region, Sacramento, CA. FINAL REPORT May 14, 2018, 144p. (<https://www.usbr.gov/mp/bdo/six-year-acoustic-telemetry-steelhead-study.html>)
- U. S. Fish and Wildlife Service. 2003. Flow-Habitat Relationships for Steelhead and Fall, Late-Fall and Winter-Run Chinook Salmon Spawning in the Sacramento River Between Keswick Dam and Battle Creek. Pages 79pp. in, Sacramento, California.
- U.S. Fish and Wildlife Service. 2006. Relationships Between Flow Fluctuations and Redd Dewatering and Juvenile Stranding for Chinook Salmon and Steelhead in the Sacramento River Between Keswick Dam and Battle Creek. Pages 94.
- U.S. Fish and Wildlife Service. 2007. Flow-Habitat relationships for Spring-run Chinook salmon and Steelhead/Rainbow Trout spawning in Clear Creek Between Whiskeytown Dam and Clear Creek Road.
- Voss, S. D., and W. R. Poytress. 2017. Brood year 2015 juvenile salmonid production and passage indices at the Red Bluff Diversion Dam. Report of U.S. Fish and Wildlife Service to U.S. Bureau of Reclamation, Sacramento, CA.
- Voss, S. D., and W. R. Poytress. 2018. Brood year 2016 juvenile salmonid production and passage indices at the Red Bluff Diversion Dam. Report of U.S. Fish and Wildlife Service to U.S. Bureau of Reclamation, Sacramento, CA.

- Voss, S. D., and W. R. Poytress. 2019. Brood year 2017 juvenile salmonid production and passage indices at the Red Bluff Diversion Dam. Report of U.S. Fish and Wildlife Service to U.S. Bureau of Reclamation, Sacramento, CA.
- Voss, S. D., and W. R. Poytress. 2020. Brood year 2018 juvenile salmonid production and passage indices at the Red Bluff Diversion Dam. Report of U.S. Fish and Wildlife Service to U.S. Bureau of Reclamation, Sacramento, CA.
- Voss, S. D., and W. R. Poytress. 2022. 2019 Red Bluff Diversion Dam Rotary Trap Juvenile Anadromous Fish Abundance Estimates. Report of U.S. Fish and Wildlife Service to U.S. Bureau of Reclamation, Sacramento, CA.
- Whitton, K. S., J. M. Newton, D. J. Colby, and M. R. Brown. 2006. Juvenile salmonid monitoring in Battle Creek, California, from September 1998 to February 2001. USFWS Data Summary Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., J. M. Newton, and M. R. Brown. 2007. Juvenile salmonid monitoring in Battle Creek, California, July 2001 through September 2002. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., J. M. Newton, and M. R. Brown. 2007. Juvenile salmonid monitoring in Battle Creek, California, October 2002 through September 2003. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., J. M. Newton, and M. R. Brown. 2007. Juvenile salmonid monitoring in Battle Creek, California, October 2004 through September 2005. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., J. M. Newton, and M. R. Brown. 2007. Juvenile salmonid monitoring in Battle Creek, California, October 2003 through September 2004. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., J. M. Newton, and M. R. Brown. 2007. Juvenile salmonid monitoring in Battle Creek, California, October 2005 through September 2006. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., D. J. Colby, J. M. Newton, and M. R. Brown. 2008. Juvenile salmonid monitoring in Battle Creek, California, November 2007 through June 2008. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., D. J. Colby, J. M. Newton, and M. R. Brown. 2010. Juvenile salmonid monitoring in Battle Creek, California, November 2008 through June 2009. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.
- Whitton, K. S., D. J. Colby, J. M. Newton, and M. R. Brown. 2011. Juvenile salmonid monitoring in Battle Creek, California, November 2009 through July 2010. USFWS Report. U.S. Fish and Wildlife Service, Red Bluff Fish and Wildlife Office, Red Bluff, CA.

- Williams, J. G. 2006. Central Valley salmon: a perspective on Chinook and steelhead in the Central Valley of California. *San Francisco Estuary and Watershed Science* 4(3).
- Williams, J. G. 2010. DRERIP Delta Conceptual Model: Life History Conceptual Model for Chinook Salmon & Steelhead *Oncorhynchus tshawytscha* & *Oncorhynchus mykiss*. Department of Fish and Game Ecosystem Restoration Program.
- Williams, T. H., S. T. Lindley, B. C. Spence, and D. A. Boughton. 2011. Status review update for Pacific salmon and steelhead listed under the Endangered Species Act. National Marine Fisheries Service, Southwest Fisheries Science Center, Santa Cruz, CA.
- Williams, T. H., B. C. Spence, D. A. Boughton, R. C. Johnson, L. Crozier, N. Mantua, M. O'Farrell, and S. T. Lindley. 2016. Viability assessment for Pacific salmon and steelhead listed under the Endangered Species Act: Southwest. 2 February 2016 Report to National Marine Fisheries Service – West Coast Region from Southwest Fisheries Science Center, Fisheries Ecology Division 110 Shaffer Road, Santa Cruz, California 95060.
- Yuba County Water Agency. 2014. Draft Biological Assessment for Central Valley spring-run Chinook salmon, Central Valley steelhead, and North American green sturgeon and draft essential fish habitat assessment. Application for new license, major project – existing dam. Volume IV: exhibit E. Yuba River Development Project, FERC No. 2246. April 2014.
- Zaugg, W. S., B. L. Adams, and L. R. McLain. 1972. Steelhead Migration - Potential Temperature Effects as Indicated by Gill Adenosine-Triphosphatase Activities. *Science* 176(4033):415–416.

5.3 Models

Models support testing alternative operations and predicting environmental responses. The following models were available to Reclamation and relevant to addressing management questions:

5.3.1 Water Operations

CalSim II is a generalized reservoir-river basin simulation model that allows for specification and achievement of user-specified allocation targets, or goals (Draper et al. 2004). CalSim II represents the best available planning model for CVP and State Water Project (SWP) system operations and has been used in previous system-wide evaluations of CVP and SWP operations (Bureau of Reclamation 2015). Reclamation and the California Department of Water Resources are advancing CalSim 3, but the model was not ready for these purposes.

5.3.2 Water Temperature

HEC_5Q is a reservoir routing and temperature model. Over the past 15 years, various temperature models were developed to simulate temperature conditions on the rivers affected by CVP and SWP operations (e.g., Sacramento River Water Quality Model [SRWQM] San Joaquin River HEC_5Q model) (Bureau of Reclamation 2008). Recently, these models were compiled and updated into a single modeling package referred to here as the HEC-5Q model. Further updates were performed under the long-term operation (LTO) EIS modeling that included improved meteorological data and subsequent validation of the Sacramento and Stanislaus River models, implementation of the Folsom Temperature Control Devices and low-level outlet, implementation of the Trinity auxiliary outlet, improved temperature targeting for Shasta and Folsom Dams, as well as improved documentation and streamlining of the models and improved integration with the CalSim II model (Bureau of Reclamation 2015). A summary of previous model calibration and validation details can be found at the following link: [DWR-1084 RMA 2003 SRWQM.pdf \(ca.gov\)](#). Reclamation is developing an updated water temperature modeling platform, but the model is not yet available for broad use.

6. Lines of Evidence

Analysis of the LTO relies on multiple lines of evidence from datasets, literature, and models.

Lines of Evidence section is currently under development and will be provided for the Public Draft Environmental Impact Statement).

This page intentionally left blank

7. Uncertainty

Future studies of high value that may benefit from special studies include estimating the juvenile production of steelhead in the Stanislaus River to evaluate the effect flow and temperature operations has on this populations. The special study considering this is:

Steelhead JPE and OMR Management

This page intentionally left blank

8. References

Literature referenced for Stanislaus SRP are listed in Section 3 above. Additional references cited or used for informational material in the document are included below.

Appendix A. Fall Pulse Flow Advice.

Appendix B. WIF Advice.

Appendix C. Spring Pulse Flow Advice.

CalFish. 2019. CalFish – A California cooperative anadromous fish and habitat data program. Middle Sacramento River salmon and Steelhead monitoring. Available: <https://www.calfish.org/ProgramsData/ConservationandManagement/CentralValleyMonitoring/SacramentoValleyTributaryMonitoring/MiddleSacramentoRiverSalmonandSteelheadMonitoring.aspx>.

Draper et al. 2004

Fortier, R., J. Nelson, R. Bellmer, and R. Nielson. 2014. Implementation Plan for the Central Valley Steelhead Monitoring Program. California Department of Fish and Wildlife, Fisheries Branch. July 2014.

National Marine Fisheries Service (NMFS). 2009. Biological Opinion and Conference Opinion on the Long-term Operations of the Central Valley Project and State Water Project. National Oceanic and Atmospheric Administration, National Marine Fisheries Service. June 2009.

National Marine Fisheries Service. 2014. Final recovery plan for the evolutionarily significant units of Sacramento River winter-run Chinook salmon and Central Valley spring-run Chinook salmon and the distinct population segment of California Central Valley steelhead. National Oceanic and Atmospheric Administration, National Marine Fisheries Service. July 2014.

Sturrock, A. M., Carlson, S. M., Wikert, J. D., Heyne, T., Nusslé, S., Merz, J. E., Sturrock, H. J. W, Johnson, and R. C. Johnson. 2020. Unnatural selection of salmon life histories in a modified riverscape. *Global Change Biology* 26:1235-1247. <https://doi.org/10.1111/gcb.14896>.

U.S. Bureau of Reclamation 2008

U.S. Bureau of Reclamation 2015