

The Science of Non-native Species in a Dynamic Delta

Delta Independent Science Board

Public Draft Report (July 14, 2020)

All links in this report have been created with meaningful text. The Uniform Resource Locator (URL) is also published to be available as a resource for those who print the document. The URL addresses that are spelled out in the document are not active links to avoid the confusion of presenting duplicate links.

This document makes use of italics and bold text for emphasis.

If you need assistance interpreting the content of this document or would like to provide public comments, please e-mail disb@deltacouncil.ca.gov.

Table of Contents

Introduction and Rationale	3
The Context of Non-Native Species in a Dynamic Delta	4
The Invasion Process	6
Findings	6
Background and Definitions	7
Ingredients for Establishment of a Non-native Species	9
Non-native Impacts on Ecosystems	11
Non-Natives in the Sacramento-San Joaquin Delta	13
Findings	13
History and Status	14
Management and Coordination	17
Recommendations	19
Individual Invasive Species: Prevention and Management	20
Findings	20
The Overall Process and Scientific Needs	21
Threat Assessment and Prevention	22
Rapid Response and Eradication	25
Control and Adaptation	26
Recommendations	27
Non-native Species in the Context of Ecosystem Management	29
Findings	29
Ecosystem Management and Non-native Species	30
The Dynamic Delta and Non-native Species	32
Recommendations	35
Ecological Restoration and Adaptive Management	36

PUBLIC DRAFT (7/14/2020)

Findings _____	36
Restoration in the context of Non-native Species _____	36
Recommendations _____	39
Management Coordination and Integration _____	39
Conclusions _____	41
APPENDIX A: Examples of Significant Non-native Species in the Delta ____	43
Bivalves and their effects on the pelagic food web _____	43
Aquatic plants _____	44
Wetland vegetation _____	48
Fish _____	49
Non-native fish and submerged aquatic vegetation (SAV) _____	52
Mammals _____	53
APPENDIX B: Panelists _____	54
References _____	54

PUBLIC DRAFT (7/14/2020)

Introduction and Rationale

The Sacramento-San Joaquin Delta Reform Act of 2009 stipulates that the Delta Independent Science Board “shall provide oversight of the scientific research, monitoring, and assessment programs that support adaptive management of the Delta through periodic reviews.”¹ Management of the Delta to realize the coequal goals stated in the Delta Reform Act is directed by the Delta Plan. The Act also stipulates that the Delta Plan should restore a healthy ecosystem by promoting “self-sustaining, diverse populations of native and valued species by reducing the risk of take and harm from invasive species.”²

The purpose of this review is to assess the state, quality, and potential usefulness of **scientific information** that helps agencies understand and manage the threats and consequences of non-native species (plants and animals) into Delta lands and waters. The overall goal of this review is to offer recommendations to improve the science needed to manage for non-native and non-native invasive species in the Delta. This review process included an extensive literature review and two panels each comprised of five experts who explored the status of science relative to non-native species in the Sacramento-San Joaquin Delta. Additionally, Delta Independent Science Board members participated in several workshops, presentations, and discussions with managers. We chose this topic because the Bay-Delta ecosystem is one of the most invaded ecosystems on the globe and reducing the impact of non-native species and protecting native species is one of the five core strategies discussed in the Delta Plan’s Chapter 4 amendment (“Protect, Restore and Enhance the Delta Ecosystem”).

Overall, there has been a tremendous amount of research done on non-native species in the Sacramento-San Joaquin Delta (hereafter, “Delta”). It is beyond

¹ California Water Code Section 85280.

² California Water Code Section 85302(e)(3).

PUBLIC DRAFT (7/14/2020)

the scope of this review to summarize all of that information or to list all of the project- or species- or technology-specific science or monitoring that has or should be done. Rather, we focus our findings and recommendations on a higher-level approach to provide managers with a scientific-based prioritization framework to make decisions. We use examples for the Delta to support our findings. The consequences of biological invasions, however, are a global concern (e.g. Pysek et al. 2020) and the scientific needs to better prevent, control, and ultimately manage invasive species are similar across ecosystems.

We begin the review by providing a broad context for considering non-native species in a dynamic Delta. We then define terms and discuss the invasive process. We provide an overview of the individual species approach to invasive-species management and prevention and discuss how non-natives fit into the broader picture of changes in species composition in a dynamic and rapidly changing ecosystem. We illustrate how science can inform management decisions at different stages in the invasive process. We follow by considering how ecological restoration may affect and be affected by non-native species, and how the continual threats from non-native species affect and are affected by the practice of adaptive management. We highlight areas in which scientific knowledge or its application in the Delta relative to the influx of non-native species could be expanded and better coordinated. Throughout, we offer recommendations to strengthen the management of dealing with non-native species in the Delta.

The Context of Non-Native Species in a Dynamic Delta

To understand, anticipate, and manage non-native invasive species in the Delta, one must consider them in the context of a dynamic and ever-changing environment. Two realities cannot be ignored.

PUBLIC DRAFT (7/14/2020)

First, today's Delta is not a pristine ecosystem. Far from it—it is one of the most heavily modified estuaries on Earth. Well before the arrival of European settlers, Native Americans altered the mosaic of species in the Delta by tending local plant species that bore acorns, fruits, and construction materials and by moving them into new locations (Zedler and Stevens 2018). Beginning with European colonization of the Americas, people mixed species between the eastern and western hemispheres (Mann 2011), a practice that has continued through to the economic globalization of today. The massive alterations that began in the mid-nineteenth century and the subsequent engineering of the Delta to support agriculture and manage water have accelerated invasions by non-native species.³ Many non-native species have become “naturalized” members of Delta ecosystems.

Second, the major forces now driving environmental change in the Delta—climate change, sea-level rise, and human uses of land and water resources—are subject to a complex interplay of global, regional, and local influences, many of which are beyond direct management. As these forces mount, environmental changes are becoming more rapid, extreme events such as droughts or deluges are becoming more frequent and more extreme, and tipping points of ecosystem change are more likely to be passed—the pelagic organism decline (POD) that occurred in the Delta in 2002 is an example (MacNally et al. 2010). The environmental turmoil created by these forces of change will provide new opportunities for non-native invaders and challenge the capacity of native species to adapt, of scientists to understand and predict ecosystem dynamics, and of managers to shepherd their land and water resources responsibly.

Everything in this report should be considered in the context of these two overarching realities. Our overall recommendation is to **encourage a broader,**

³ Whipple et al. (2012) and SFEI-ASC (2014) review the history and current status of Delta landscapes and ecosystems.

PUBLIC DRAFT (7/14/2020)

more forward-looking, integrated approach to non-native species science in the Delta to inform management goals.

The Invasion Process

Findings

- The definition of an **invasive species** is a non-native species that does or is likely to cause environmental or economic harm or harm to human health. It is based on a human value judgement.
- All invasive species are non-native, but not all non-native species are invasive.
- In general, managers have favored native over non-native species to conserve biodiversity, ecosystem services, and historical Native American cultural functions.
- Non-native species are one of the five fundamental drivers of ecosystem change.
- Non-native species can disrupt food-webs, nutrient and contaminant cycling, habitat structure, and ecosystem services.
- Management can help buffer an ecosystem from future invasions by concentrating on the two processes that humans can control: reducing or eliminating invasion pathways and reducing ecosystem vulnerability to new invaders.
- Once a new non-native species has become established in an ecosystem, the structure, composition, and likely the functioning of the ecosystem are changed.
- The species pool in an ecosystem is dynamic, leading to a continual reshuffling of native and non-native species.

PUBLIC DRAFT (7/14/2020)

Background and Definitions

Globally, invasive species are one of the greatest threats to the ecological integrity of ecosystems (IPBES 2019) and may have contributed to 25% of the global plant extinctions and 33% of the animal extinctions (Pysek et al. 2020). The emergence of invasion ecology as an area of broad scientific and public concern dates from the publication of Charles Elton's book, *The Ecology of Invasions by Animals and Plants* (1958). Elton cast the challenge of invasive species using a military metaphor:

"I have described some of the successful invaders establishing themselves in a new land or sea, as a war correspondent might write a series of dispatches recounting the quiet infiltration of commando forces, the surprise attacks, the successive waves of later reinforcements after the first spearhead fails to get a foothold, attack and counter attack, and the eventual expansion and occupation of territory from which they are unlikely to be ousted again" (Elton 1958: 109).

Although this militaristic metaphor may no longer be appropriate (Davis et al. 2011, Janovsky and Larson 2019), it does capture many of the features of the battle against invasive species and their characterization as harmful. Invasive species are considered to be one of the five direct drivers of ecosystem change along with climate change, resource use, habitat alteration (land use) and pollution (Millennium Report, 2005). Accordingly, the literature on this topic is extensive.

The concepts of invasive and non-native species and related terms have been controversial since their beginnings; different researchers adopt different definitions for their own reasons, and the controversies will continue as environmental change quickens. The terminology for non-native species is also confusing, confounded and inconsistent (e.g. Shrader-Frechette 2001, Colautti and MacIsaac 2004). A variety of terms has been used to denote a non-native

PUBLIC DRAFT (7/14/2020)

species, including alien, nonindigenous, exotic, invader, weed, aquatic nuisance species, introduced species, and foreign species. The definitions are more well defined in legislation and executive orders. A **non-native species** is a species that is not originally from the ecosystem in which it now occurs. Using the National Invasive Species Management Plan (Beck et al. 2006), we use the definition of an **invasive species** as “a species that is non-native to the ecosystem under consideration and whose introduction causes or is likely to cause economic or environmental harm or harm to human health.” The National Invasive Species Council added that invasive species are those introduced to an area as a result of intentional or unintentional human actions. In general, managers have favored native over non-native species to conserve biodiversity, ecosystem services, and historical Native American cultural functions.

By this definition, all invasive species are non-native, but not all non-native species are invasive. The two essential elements in the definition of an invader are that the species is non-native and that it causes harm. Whether a non-native species entering an ecosystem causes harm is a matter of human values, which can change or differ among groups of people. A non-native species can become invasive as its impact changes or becomes better known and some species can be considered both detrimental and beneficial. For example, sport fishers in the Delta currently value non-native striped bass (*Morone saxatilis*) that were introduced and became established over a century ago, whereas others emphasize the harm the bass now may cause by preying on native fishes (Moyle 2011; 2020). Striped bass are now managed as a recreational resource in the Delta. Therefore, determining whether a species should be labeled “invasive” can depend on how people perceive the economic and environmental benefits and costs of the species and how these are balanced (Beck et al. 2006), and different people do it differently.

For management purposes, native species are generally considered to be those species present in an area when Europeans first arrived and described what they found. Pysek and Richardson (2010) suggest that native species “evolved in a

PUBLIC DRAFT (7/14/2020)

given area without human involvement or ... arrived there by natural means ... from an area in which they are native.” Thus, species such as cattle egrets (*Bubulcus ibis*), which emigrated from their native Africa on their own and colonized much of the Americas, are not generally considered invasive. By this measure, a human vector must be involved for a species to be called invasive.

As more species expand their ranges from areas in which they are native into new areas in response to climate changes, however, determining whether a species is or is not native may be less important than determining whether it meets the second defining element: causing harm. For example, barred owls (*Strix varia*), native to eastern North America, have expanded into forests of the Pacific Northwest where they were historically not present. They compete with federally threatened northern spotted owls (*Strix occidentalis caurina*), displacing them from many areas and hastening their decline (J.D. Wiens et al. 2014). Should barred owls be considered an invasive species?

Ingredients for Establishment of a Non-native Species

There are three essential ingredients for the successful establishment of a non-native species in an ecosystem.

- 1) There must be a **pathway or corridor** that allows the species to traverse the natural barriers that may prevent the species from getting to an ecosystem. These barriers can simply be the distance or the presence of inhospitable habitats. There are natural ways to break through these barriers that vary from continual range expansion to changes in intervening habitat to accidental transport by another organism (e.g., aquatic organisms attaching to water birds). The success of establishment of a non-native species is often dependent on the number of introduction events and the number of individuals introduced (see review by Pysek et al. 2020).

PUBLIC DRAFT (7/14/2020)

Human activity has created multiple pathways for invasions through deliberate release with or without intent (stocking, bait release), hitchhikers on commodities (e.g. insects) or on transport vectors (e.g. biofouling, ballast water, boats), escape from captivity (aquaria pets), or creation of anthropogenic pathways (e.g. canals and water diversions).

- 2) The second essential ingredient is **a match of the physical, biological, or chemical habitat requirements of the potential non-native species to those of the receiving ecosystem**. Are habitat conditions suitable for growth, reproduction and persistence of the non-native species in this ecosystem or do predators, competitors, or adverse habitat conditions restrict establishment of the new species? In essence, the basic drivers of ecosystem change driven by climate, habitat alterations, pollution, extreme events, and resource use can and will change the habitat receptivity to different types of non-native species.

Humans have altered the receiving habitats and therefore have altered their susceptibility to invasion by different non-native species. Human alterations can include changes in hydrological flow amounts and patterns, habitat structure, species composition (resource exploitation), nutrient and pollution input, food-web disruption, and even the initial influx of non-natives that can change habitat vulnerability to additional non-native species.

Given the above, then prevention of new invaders should be focused on reducing ecosystem vulnerability and pathway restrictions.

- 3) The third ingredient for the successful establishment of a non-native species is often related to the inherent biological and ecological traits of the individual species—**the habitat and reproductive requirements and abilities of the potential invasive species**. Some species are better adapted to expand into new environments because they are generalist feeders, have rapid reproductive capabilities, have high tolerance for a

PUBLIC DRAFT (7/14/2020)

wide range of environmental conditions, or have greater resistance to predators. Ultimately, the success or failure of a species that enters an ecosystem will depend, in part, on these characteristics, although the relationship are challenging to define quantitatively (e.g. Ricciardi and Rasmussen, Kolar and Lodge 2001, Marchetti et al. 2004).

Non-native Impacts on Ecosystems

Once a new non-native species has become established in an ecosystem, the structure, composition, and, likely, the functioning of the ecosystem are changed. To evaluate the science underpinning efforts to address non-native species problems in the Delta, invasion can be considered as one aspect of the broader dynamics of the community of species occurring in the Delta (the “species pool”; Figure 1).

Several forces drive changes in the species pool. These drivers—climate change, sea-level rise, land-use change, habitat alteration, hydrological changes, resource use, pollution and nutrient loading, droughts, and a host of other human actions—all affect species and their habitats directly and indirectly. As a consequence, the species pool in an area of interest is in a continual state of flux, with changing population levels of species already present, additions of new species from elsewhere, and loss of species previously present in the pool. Additions come from immigration of species moving of their own accord, intentional human introductions of new species, (e.g., assisted migration or stocking), or accidental or careless introduction through human-facilitated pathways (e.g., release of bait fish, clams hitchhiking on recreational boats).

Whether a newly arriving species becomes established depends on abiotic conditions, the characteristics of species that moved into the area earlier, and how they assembled themselves into ecosystems. Once established, a non-native species may affect the persistence or decline of species already present and those that arrive subsequently. Losses of species from the pool occur when a species becomes extinct or is extirpated from the area of interest or when a

PUBLIC DRAFT (7/14/2020)

species disappears because individuals and population centers have moved elsewhere (e.g., as a result of climate change). There are also transients in the species pool such as migratory birds and fishes. The species pool of any location therefore contains a mixture of native and non-native species that changes over time, creating an ever-changing mosaic of ecosystems over a broader area as species move among locations.




Figure 1. A conceptual model of changes in the species composition (the “species pool”) of an ecosystem, leading to multiple consequences.

Compositional changes in the species pool can have a variety of ecological, economic, or sociological consequences (Figure 1). Ecologically, altered competitive or predator-prey relationships among species may disrupt food webs. The effects on native species that are rare or declining in abundance may be especially great, leading some to be extirpated. If these species are legally recognized as threatened or endangered, there will be political and economic as well as ecological consequences.

PUBLIC DRAFT (7/14/2020)

Other consequences of changes in the species pool may affect human interests more directly. Ecosystem services provided by existing species and biological communities may change. For example, new species may alter the biological, hydrological, or physical structure of the ecosystem (e.g., nutria, *Myocastor coypus*, burrow into levees). Changes in the composition of aquatic vegetation, such as the recent dominance of the Delta by dense growths of Brazilian waterweed, can alter water flows, temperature, and chemistry and can affect other elements of aquatic ecosystems as well as the quality and quantity of water available to people.

Changes in the composition of a species pool and their consequences, of course, are just changes. It is *people* who determine whether the individual or collective changes are good, bad, or benign, depending on how they affect something about the system that people value, for whatever reasons. In some instances, the introduction of a new species into an area may have little observed effect on other species, ecosystem processes, or how humans use or manage the system (Matern and Brown, 2005). In other situations, a new species may be valuable to people (e.g., striped bass) or increase or alter the productivity of food webs (e.g., Liao et al. 2018). When new species have harmful consequences, they are labeled invasive species.

Non-Natives in the Sacramento-San Joaquin Delta

Findings

- The Sacramento-San Joaquin Delta is one of the most invaded estuaries in the world.
- Reducing the impact of non-native species and protecting native species is a core strategy of the Delta Plan.
- Several factors have facilitated Delta invasions, including ballast-water pathways through the San Francisco Bay and severe habitat restructuring for land and water use.

PUBLIC DRAFT (7/14/2020)

- The vulnerability of disturbed environments to invasion is well documented in other ecosystems and has been substantiated by studies in the Delta.
- Changes in the Delta over the past decades have generally favored non-native species (fish, at least) at the expense of native species.
- Science dealing with non-native species in the Delta has been extensive.
- Impacts of invaders on the Delta Ecosystem have been large but attributing specific impacts to specific species is challenging scientifically because science is reactive (done after an invasion) and data are limited.

History and Status

The San Francisco Estuary (including the Delta) has been described as one of the most invaded estuaries in the world (Cohen and Carlton 1998). Because the Delta-San Francisco Bay Area is one of 25 global biodiversity hotspots of highest priority for conservation, the threat of invasive species is a major environmental concern. More than 200 non-native species have invaded the Delta's aquatic and terrestrial habitats. The many transport pathways that bring non-native species into San Francisco Bay—international shipping, recreational boating and fishing, horticulture and pet industries, agriculture, and deliberate introduction—have facilitated their movement into the Delta (Luoma et al. 2015). These pathways, combined with the Delta's highly altered landscape, have facilitated the establishment of many non-native species (Ruiz et al. 2011). About one quarter of non-native species introduced to the estuary are arthropods, followed by mollusks, fish, and vascular plants (Cohen and Carlton 1998).

Well before the arrival of non-European settlers in the Delta, Native Americans altered the mosaic of species by tending local plant species that bore acorns, fruits, and construction materials and by moving them into new locations (Zedler and Stevens 2018). Subsequent people introduced domesticated grazers (horses, cattle). Grasses were favored by grazing and by fires set by lightning and by Native Americans. With the settling of European immigrants, California's

PUBLIC DRAFT (7/14/2020)

Central Valley was gradually converted from native to the Delta alien grasses, and the Delta was engineered to support agriculture.

Introductions began to accelerate as ships started entering San Francisco Bay in 1775. As global shipping into the Bay increased around 1850, introduction pressure intensified (Cohen and Carlton 1995, Ruiz et al. 2000). Introduction rates have increased since the mid-1900s; about half of non-native species recorded in 1995 were introduced after 1960 (Cohen and Carlton 1998). This increase coincides with a time of growing international commerce from East Asia, the opening of new ports in the 1970s, faster ships, and increasing anthropogenic disturbance (Carlton et al. 1990, Carlton 1996). In particular, habitats were altered by increasing hydrological management through freshwater diversions beginning in the 1920s and major dam construction on the Sacramento River and its tributaries between 1945 and 1968 (Arthur et al. 1996, Winder and Jassby 2011). Conditions will continue to change into the future (Lund et al. 2010). Salinity will change in different parts of the Delta with changes in hydrological regimes (Fleenor et al. 2008), with cascading effects on Delta ecosystems and fish (Moyle and Bennett 2008). These transformations of the Delta facilitate the establishment and persistence of new species.

The vulnerability of disturbed environments to invasion is well documented in other ecosystems and has been substantiated by studies in the Delta (Leidy and Fiedler 1985, Feyrer and Healey 2003, Conrad et al. 2016). Hydrologic alterations—especially water diversions, altered flows, and increased water temperatures—have exacerbated drought-like conditions, which are linked to increasing invasions by non-native zooplankton that have in turn created conditions more favorable to non-native fish (Feyrer and Healey 2003, Winder et al. 2011).

Appendix A summarizes some specific examples of the impacts of non-natives in the Delta. Non-native species can often outcompete, prey upon, and exclude native species. The continuous arrival and spread of non-natives have displaced

PUBLIC DRAFT (7/14/2020)

native aquatic vegetation, decimated native fish populations, contributed to the decline of native biodiversity, altered food webs and ecosystems, structurally damaged both natural and constructed habitats, and affected ecosystem services such as the provision of clean water (Simberloff and Rejmanek 2011). As in other estuaries, the Delta ecosystem also is vulnerable to invasion because brackish waters generally have fewer indigenous animal species than other habitats, facilitating the establishment of non-native species (Cohen and Carlton 1998, Wolff 1998, Cloern and Jassby 2012). The range of salinity conditions exposes the Delta to potential invasion by non-native species through a multitude of vectors and creates conditions favoring establishment once they arrive.

Some introduced species have had more substantial environmental and economic impacts than others due to their capacity to reshape their environment, with cascading effects on habitat, nutrient and contaminant cycling, and trophic structure (Kimmerer et al. 1994, Crooks 2002, Sousa et al. 2009). Significant habitat-altering invasive species include several species of aquatic plants that alter flows and create novel habitat for non-native fish (Brown and Michniuk 2007, Loomis 2019). Filter-feeding bivalves have altered benthic and pelagic food-web structure and nutrient cycling. Sometimes species exhibit a boom and bust invasion in which abundances and impacts can change significantly, as with the Chinese mitten crab (Box 1).

PUBLIC DRAFT (7/14/2020)

Box 1. The Chinese Mitten Crab: A Boom and Bust Invasive in the Bay and Delta

Chinese mitten crabs are medium-sized crabs named for their hairy, mitten-like claws (Rudnick et al. 2005). They are native to coastal rivers and estuaries of central Asia and have invaded several European countries over the past century. Discovered in South San Francisco Bay in 1992, the mitten crab spread rapidly to cover several thousand squared kilometers surrounding the Bay and Delta (Rudnick et al. 2000). Introductions may have occurred through ballast-water discharges, although there was initial speculation that it was purposeful because of the value of their roe.

Chinese mitten crabs are catadromous, which are species that live in freshwater but migrate to more saline habitats to breed, and are associated with tidally influenced portions of Bay tributaries as young juveniles; with freshwater streams ≤ 250 km from their confluence with the Bay) as older, migrating juveniles; and with the open waters of the Bay as reproductive adults after migrating from fresh water to reproduce between late fall and early spring (Rudnick et al. 2000, 2003). Chinese mitten crabs have been of widespread environmental concern because of their extreme abundance and burrowing behavior, which causes bank erosion. Between 1995 and 2001, burrow densities increased five-fold in tidal portions of the banks in South Bay tributaries (from a mean of 6 burrows per meters squared in 1995 to >30 burrows per meter squared in 1999). Population size peaked in 1998, with 750,000 crabs counted in fall migration in a North Bay tributary. Abundance subsequently declined greatly; 2,500 crabs were counted in the same river system in 2001 (Rudnick et al. 2003). They are rarely encountered in the Bay and Delta today.

Chinese mitten crabs are also of concern because they accumulate higher concentrations of mercury than crustaceans living in the water column (Hui et al. 2005). Because their predators include fish, birds, mammals, and humans, their mercury burdens have an exceptional potential to impact the ecosystem and public health. Chinese mitten crabs also damage nets used in commercial fisheries (Rudnick and Resh 2002).

Management and Coordination

Given the prevalence of non-native species in the Delta, the Delta Plan identifies reducing the impact of non-native species and protecting native species as a core strategy in the Ecosystem Goal (Box 2). Several interagency programs have also been formed to prevent, detect, and manage invasive species, including the

PUBLIC DRAFT (7/14/2020)

Delta Interagency Invasive Species Coordination Team, which is organized by the Sacramento-San Joaquin Delta Conservancy and aims to strengthen coordination among agencies to detect, prevent, and manage invasive species.

Box 2. Reducing Impact of Non-native Species is a core strategy in the Delta Plan.

Reducing the impact of non-native species and protecting native species is one of the five core strategies discussed in the Delta Plan's Chapter 4 amendment ("Protect, Restore and Enhance the Delta Ecosystem"). Within this strategy, the Plan recommends that state and federal agencies should prioritize and implement actions to control non-native species (ER R7), including communication and funding for a rapid response to invasive species (Delta Stewardship Council 2020). The Plan classifies non-native species into four categories: naturalized species, widespread and unmanaged species, widespread and managed species, and emerging species of concern. Invasive species are described as non-natives whose introduction may cause harm to the economy, environment, or human health.

The Plan addresses the specific threats posed by several invasive species, including aquatic weeds (water hyacinth, Brazilian waterweed, water pennywort, Eurasian water milfoil, and parrot feather), overbite clams, and zooplankton. In addition, it explains the potential threat of invasions by zebra mussels, quagga mussels, and nutria. The Plan also discusses measures and entities that have been established to prevent introduction of non-native species. California law requires that ships entering from outside the United States Exclusive Economic Zone either retain, properly exchange, or discharge ballast water to a treatment facility to reduce the chances of introduction. In addition, the California State Lands Commission limits the allowable concentration of living organisms in discharged ballast water.

The Delta Plan recommends increasing funding and communication among agencies for invasive species management.

The California Invasive Plant Council is a non-profit organization that catalogs invasive plants present in California, and the California Department of Food and Agriculture has lead authority to control of noxious weeds in California. In addition, the Delta Region Areawide Aquatic Weed Project is a collaboration among academic and governmental agencies tasked with sustainably managing

PUBLIC DRAFT (7/14/2020)

aquatic weeds in the Delta. More broadly, the Invasive Species Council of California ([ISCC website](http://www.iscc.ca.gov/), <http://www.iscc.ca.gov/>) aims to coordinate and strengthen the various organizations that address invasive species in the state of California.

Recommendations

The science dealing with non-native species in the Delta is extensive and has largely emphasized: 1) prevention, early detection and rapid response, eradication, assessment and monitoring, and control of individual species (e.g., nutria) or groups of similar invaders (e.g., emergent aquatic vegetation); 2) retrospective impact assessment (e.g., the effects of invasive clams); and 3) development of new technologies for monitoring (e.g., remote sensing and eDNA) (e.g. Jerde et al. 2011, Baerwald et al. 2012, Stoeckle 2016, Hosler 2017, Khanna et al. 2018b). See Appendix A.

We recognize that there are many scientific needs at the project level, species level, monitoring level, or technology level. These span topics such as the development of safe control measures (e.g. herbicides), development of new monitoring tools (eDNA, remote sensing), and evaluation of pairwise species relationships (e.g. striped bass and delta smelt) to more challenging questions like better defining the role of an individual invader (e.g. *Corbicula*) in nutrient cycling. Clearly, all of these types of projects are important but need better prioritization. **We recommend that a series of workshops or syntheses be conducted to develop a detailed set of Science Priorities for dealing with invasive species that defines short-term and long-term science needs.** Recent workshops like the 2019 assessment of remote sensing technology and status for invasive aquatic vegetation (references) are good examples of the type of approach that is needed.

One of the primary impacts of non-native species is to disrupt or change the food web and nutrient cycling. Understanding the role of non-native species (potential, existing, or outgoing) in the food web is fundamental for predicting and evaluating

PUBLIC DRAFT (7/14/2020)

impacts (David et al. 2017). **We recommend the development and testing of a comprehensive food-web model for the Delta.** This type of model is most effective for policy if it is spatially-explicit, can be driven by changing environmental conditions, and is open source (e.g. DeMutsert et al. 2018, Schuckel et al. 2018). Several shelf-ready models already exist (e.g. ECOSIM/ECOPATH/ECOSPACE, e.g. Vassslide et al. 2016) but we believe a coordinated effort to evaluate the most appropriate approach for the Delta is needed (e.g. Schuckel et al. 2018).

Individual Invasive Species: Prevention and Management

Findings

- The primary goal of management in the Delta is to prevent the introduction of potential invaders to the ecosystem. Decisions are thus mostly focused on the different phases of individual invaders: threat assessment, early detection and rapid response to eradicate, control, and (if all else fails) adaptation.
- A prioritized list of potential invaders can be built for the Delta through a robust threat assessment.
- It is up to management to decide action levels. At what level of risk does one decide whether to take action take no action.
- Science and management are clearly linked and must be integrated. Each management goal/action requires science.
- Identification of shared invasion pathways for multiple species can enhance prevention efforts.

PUBLIC DRAFT (7/14/2020)

- Monitoring is essential to assess the effectiveness of prevention, to detect new non-natives, and to map the spread and abundance of established non-natives.
- Rapid response for eradication or control requires resources and agency commitment and coordination.




Figure 2. Stages of management and responses in dealing with a potential invasive species. Ideally, all of the stages and responses are informed by science and monitoring informs action.

The Overall Process and Scientific Needs

The science of non-native species includes both the science associated with individual species and the ecosystem-level science that addresses ecosystem services and function.

The general management protocol for dealing with individual invasive species is well established at local, state, national, and local levels. The responses range from prevention, early detection, and rapid response to eradicate individual

PUBLIC DRAFT (7/14/2020)

species at the early stages to the control or eventual adaptation to dealing with a well-established invader if all else fails (Figure 2). Each stage in the management decision process requires scientific information.

Threat Assessment and Prevention

Ultimately, the primary goal of management is to prevent the introduction of potential new invaders to the ecosystem. Efforts are usually targeted at primary pathways for invasions and at species anticipated to have significant impacts. A prioritized list of potential invaders is critical for setting prevention and detection goals and for managing public expectations. This list can be built through a robust threat assessment.

The first step is to conduct a **threat assessment** for the species (Figure 2). Science should be used to assess risks and identify species that have a high probability of entering the ecosystem of interest, becoming established, and doing harm (Srebalienė et al. 2019). Elements of a scientific risk assessment should include: 1) an assessment of the ability of the potential invader to thrive in the new ecosystem, which might include a comparison of the habitat requirements of the potential invader (e.g. including growth and reproductive potential, food and habitat availability, and risk of predation) relative to the habitat characteristics of the ecosystem; 2) an evaluation of the potential and realistic pathways of entry (e.g. how porous are the boundaries of the ecosystem to this particular species?); and 3) a measure of the degree of harm from a successful invasion.

Several tools are available to assess the risks and impacts of potential invaders. Over 70 tools were identified in a review of the topic by Srebalienė et al. (2019). The principal aim of these tools is to identify and prioritize the major species of concern and the major pathways so that prevention techniques can be employed and monitoring can be established to detect their presence. Forecasting a new invader requires a comparison of the habitat requirements of the potential

PUBLIC DRAFT (7/14/2020)

invader with the habitat characteristics of the receiving ecosystem. A further consideration is evaluating the spread potential of the individual species.

Gauging the harmful or beneficial impacts of an invading species is a judgment that can draw on a variety of quantitative and qualitative tools. These can range from expert opinion and ratings (developed for the State of California by the [California Invasive Species Committee \(ISCC\)](http://www.iscc.ca.gov): <http://www.iscc.ca.gov>); to observations of the species in nearby or similar habitats (e.g., zebra and quagga mussels, nutria), to a more scientific and quantitative approach including comparison of the species' habitat requirements to habitat availability in the ecosystem of interest, to risk-based decision models (e.g., Wu et al. 2010). For example, the ISCC was asked to create a list of "invasive species that have a reasonable likelihood of entering or have entered California for which an exclusion, detection, eradication, control or management action by the state might be taken" (CISAC Charter, Article IIIB)." Expert opinion and comments were used (in 2010) to rate individual species (scale of 1 to 5) on criteria such as spreading rate and amount; damage or benefit to culture, health, ecology, agriculture, and infrastructure; and the state's ability to detect and control. We could not find a similar list for the California Delta. The California Department of Fish and Wildlife has also listed 21 species⁴ and has active (mainly educational) programs that strive to prevent these species from invading additional wildlands and waterways.

Science can define the risk, levels but it is up to management to decide action levels. How high does the risk need to be (and in what units) to trigger a response or how low does the threat need to be to essentially take no action? How does one balance the threat of a species that has a high probability of entering the ecosystem but low human impact against a species that can cause extreme harm or damage but has a low probability of introduction?

⁴ [California Department of Fish and Wildlife website on invasive species:](https://wildlife.ca.gov/Conservation/Invasives/Species)
<https://wildlife.ca.gov/Conservation/Invasives/Species>

PUBLIC DRAFT (7/14/2020)

Once a species has been identified as a threat, the next management step is prevention (Figure 2). Prevention is usually targeted at eliminating the primary pathway(s) for the species to enter the ecosystem. Science is needed to identify the likely pathways and the most effective methodology to restrict that pathway for the target species. In some cases, this might be done through an approach targeted on specific species, such as inspecting boats traveling into a region or a particular ecosystem.

One of the best national examples of threat assessment and pathway interdiction involves zebra and quagga mussels (*Dreissena polymorpha* and *Dreissena bugensis*). These mussels entered the Great Lakes via ballast water and have had ecosystem-level impacts on water quality, fisheries production, and even water supply and power intakes. The economic cost has been large.⁵ The species have spread throughout much of the country (see references). Studies have focused on predicting the potential for invasions into different ecosystems by comparing the habitat requirements and restrictions of zebra mussels (based on temperature, salinity, pH, flow rates, and calcium concentrations) to potential receiving waters (e.g. Whittier et al. 2008). Other studies have developed risk-based decision models focused on potential food-web disruption and other impacts (e.g. Wu et al. 2010).

Managing pathways has ranged from boat inspections for overland transport to extensive education programs and outreach, such as the nationwide 100th Meridian Initiative.⁶ Dreissenid mussels have entered the state of California and the California Department of Fish and Wildlife has produced Guidance for Dreissenid Prevention Program.⁷

⁵ See [AIS Economic Impacts Website](http://www.aquaticnuisance.org/resources/ais-economic-impacts)[AIS Economic Impacts Website](http://www.aquaticnuisance.org/resources/ais-economic-impacts):
<http://www.aquaticnuisance.org/resources/ais-economic-impacts>

⁶ See [The 100th Meridian Initiative](https://www.fws.gov/fisheries/ANS/pdf_files/100thMeridian.pdf) website:
https://www.fws.gov/fisheries/ANS/pdf_files/100thMeridian.pdf

⁷ See [Guidance Document](https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=140345&inline):
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=140345&inline>

PUBLIC DRAFT (7/14/2020)

Pathway analyses can be effective to explore and stop the potential corridors for multiple species introductions. For the Delta, the legislation controlling ballast-water release into the San Francisco Bay is an example of controlling a key pathway. The California Marine Invasive Species Program (CMIS) was designed to reduce the risk of introducing non-native species through ballast-water discharge and was established through legislation (Ballast water management for control of nonindigenous species act of 1999, reauthorized and expanded in the Marine Invasive Species Act of 2003). These and subsequent regulations have helped to regulate ballast-water discharge and biofouling (e.g. Scianni et al. 2019).

Monitoring targeted toward individual species or as part of a more general sampling program is required to provide the data needed to map and assess the effectiveness of a prevention program. This requires knowing the potential habitats of a species and effective means to catch it and assess its abundance. Monitoring can be done on a broader scale to look for invaders (DNA analyses, taxonomic training).

Rapid Response and Eradication

If prevention fails, rapid response to gather more information (e.g. surveys) and eradicate an invader is the next management step. Eradication means that no individuals remain of the invading species. A science based, species-specific rapid-response plan is required so that effective tools can be used to eliminate a species. Rapid response protocols can be set up by a team including multiple agencies and citizen advisories that are largely independent of the particular species that may be invading.

Few invaders have actually been eradicated. Success has been greatest when invaders were detected at an early stage and in a small region. An example is *Caulerpa taxifolia*, a macroalga that has been highly invasive in the Mediterranean Sea. Prompt action was taken to eradicate the species when it was discovered in Southern California in 2000 and it was ultimately declared

PUBLIC DRAFT (7/14/2020)

eradicated in 2006 (Anderson 2004). Currently there is an integrated program in the Delta to survey the Delta and eradicate any new appearances of nutria. The California Multi-agency Response Team is collaborating to eradicate the Delta population. It began as an emergency Incident Command System in 2018 and a formal Nutria Eradication Program in 2019. The Nutria Eradication Program had caught over 1,000 nutria by May 2020 (See Appendix A).⁸

Control and Adaptation

At what point does one give up on eradication? Once an invader has become established, science is needed to assess the impact of the invader and the most effective ways to map the spread and assess, control, or limit the impact of the invasion. Controls can limit the spread, reduce the abundance, or lessen the impact of the invader. Various control techniques have been used, including manual (hand removal), mechanical (backhoe, harvester, power tools, etc.), chemical (pesticides – herbicides, fungicides, rodenticides, etc.), cultural (changing a disturbance regime to favor desirable species through grazing, controlled burning, active revegetation), biological (biocontrol agents such as bugs or pathogens), and integrated pest management (using a combination of techniques for greatest efficacy; for example, mowing weeds first to reduce biomass then spraying re-sprouts with herbicides). In the Delta, continual mapping and control of emergent vegetation is an example of the degree of effort involved (See Appendix A).

An invading species may be resistant to control efforts or the efforts may fail or come too late. Often an invader is not even detected (or recognized as causing harm) until it becomes well established (e.g. *Corbicula* in the Delta). This can happen, for example, if the invading species is small or cryptic or otherwise escapes notice until it has reached a level that allows it to persist and grow. It may take some time before a new species becomes established, its population

⁸ [California Department of Fish and Wildlife's Nutria website:](https://wildlife.ca.gov/Conservation/Invasives/Species/Nutria/Infestation)
<https://wildlife.ca.gov/Conservation/Invasives/Species/Nutria/Infestation>

PUBLIC DRAFT (7/14/2020)

expands, and it can be linked to a change in ecosystem services that we value. Such time lags and delayed impact assessments complicate management responses and require ongoing monitoring (e.g. alligator weed in the Delta).

Finally, adaptation implies that the species has established itself in the ecosystem and ecosystem-level management (or other individual species management) must consider this species as part of the ecosystem. Management must then adapt to the presence of the non-native species in the ecosystem (Figure 2). In some instances, the invader may fit into an ecosystem with minimal observable effects on other species or little disruption of ecosystem functions—it has become integrated into the ecosystem (“naturalized”) and no longer meets the definition of an invasive species (i.e., causing harm).

Recommendations

A prioritized list of those species that pose the greatest threat to the California Delta should be developed. This would allow funds to be directed at prevention, effective stakeholder engagement and education, monitoring, and early detection of those species. Such a list has not yet been developed for the Delta. Management agencies in the Delta are working within the context of large statewide and national effort but should consider the greatest new threats to the Delta.

The Delta is highly vulnerable to invasion by new aquatic species entering from San Francisco Bay or elsewhere in or beyond California. A prioritized list of potential invaders can be built through a robust threat assessment and the development of risk-based decision models (e.g., Wu et al. 2010).

The threat assessment should include the development of comprehensive quantitative models to predict potential impacts of new invaders on ecosystem structure and function, including habitat occupancy (cf. Durand et al. 2016; Tobias et al. in press). *Forecasting the impacts of a potential invader requires better understanding of food-web disruption and interactions and*

PUBLIC DRAFT (7/14/2020)

insights into predation, competition, energy and nutrient flow, and habitat structure. As mentioned before, a quantitative, spatially and temporally explicit food-web model (such as ECOSIM with ECOSPACE) for the Delta would be a good place to start.

A uniform framework for applying spatially explicit habitat models for current and potential invaders should also be developed. These can be similar to life-cycle models but be generalized so that individual species needs can be inserted. These models can be used to assess the probability of successful invasion and potential **ecological or environmental impacts**.

An analysis of the anticipated economic and environmental impacts of the highest priority new invasive species should they become established in the Delta. *This will allow actions to be further prioritized on the most harmful species, allow for enhanced stakeholder engagement and set expectations and minimize surprises to the broader community.*

Non-native Species in the Context of Ecosystem Management

Findings

- Expanding the focus of management beyond individual species allows their impacts and functional roles in ecosystems to be considered in an ecosystem context.
- After-the-fact analyses of non-native impacts are challenging because of multiple, interactive effects in the ecosystem.
- The rate, type, and impact of new introductions are intertwined with other major driving forces that change ecosystems, including resource use, climate change, pollution, and habitat alterations.
- Non-native species can have ripple effects that facilitate further invasions.
- Management of non-native species must be undertaken in the context of ecosystem dynamics; the species pool is in flux, leading to a continuing reshuffling of native and non-native species and changes in ecosystem structure, function, and services.
- Ongoing, targeted monitoring is essential; new and emerging technologies should be used.
- Changing habitats will alter the suitability of the Delta to different species and therefore change risk assessment.
- It is important to consider the consequences of anticipated climate change in the Delta in forecasting future invasions.

PUBLIC DRAFT (7/14/2020)

- With climate change, predicting invaders and the course of invasions will be more difficult and require ecological and life-history information on potential invaders.

Ecosystem Management and Non-native Species

The coequal goals of the Delta Plan call for “protecting, restoring and enhancing the Delta Ecosystem.” As mentioned earlier, Delta ecosystems, defined in part by the species pool, are undergoing continual changes.

If an invasive species becomes established, it becomes a participant in the functional processes of the ecosystem—as a competitor or predator of other species, a node in the ecosystem food web, a contributor to biogeochemical cycles, as habitat for other species, or other functional roles. Functionally, the line between a native species and an established non-native species begins to blur (Aquilar-Madrano et al. 2019). It may then be less important for managers to focus on the degree of nativeness of a species than on the functional role it plays in the ecosystem.

As many of the examples we describe in Appendix A illustrate, the roles of invasive species are often disruptive. They alter aspects of the structure, composition, and function of ecosystems that we wish to maintain. In some situations, however, the impacts of a new non-native are benign from a human perspective or do not warrant the costs of eradication, control, or ongoing management. Consequently, we must adapt to the presence of the non-native species. Determining an appropriate course of action should include an assessment of the functional role the non-native species has come to play in the ecosystem. This requires that we not only know the ecology and habitat requirements of the non-native, but that we understand the strengths of its interactions with other species, its food-web relationships, how it affects water quality and nutrient cycling or hydrological flows, and how it fits into a myriad of ecosystem processes. Unfortunately, our present knowledge of the details of how Delta ecosystems function is generally inadequate to support such assessments.

PUBLIC DRAFT (7/14/2020)

Invasive species become established in an ecosystem because conditions there fulfill their ecological niche requirements, either because the invader excludes some native species that previously occupied that niche or because there was no species present that had the same ecological niche requirements. Absence of controlling predators can also be important. Perhaps the non-native species replaces a species that became extinct centuries or millennia ago (Perino et al. 2019) or environmental changes have created new habitats (like rivers turning into calm ponds or lakes). Whether the species are functional equivalents can only be hypothesized based on structure.

Ecosystem “sustainability” or protection does not mean unchanging stasis. As the species pool changes, managers need to assess species’ functions and determine the benefits and costs of changes in dynamic ecosystems. New tools are becoming available for predicting, tracking, and controlling invaders. Dick et al. (2017) created a Relative Impact Potential metric to predict the likelihood and magnitude of ecological impacts of invasive species, using data on the numerical responses and functions derived from other populations elsewhere. Foxcraft (2009) established “thresholds of potential concern” as triggers to begin controlling non-native species in the adaptive management of South Africa’s Kruger National Park. Such approaches may help to shift the management of invasive species from response to prevention.

Godoy (2019) challenged researchers to uncover “emergent properties” of ecosystems being invaded by considering multispecies assemblages and learning how communities change once invaded. Efforts focused on just two competing species at a time (e.g., a native and non-native) miss the emergent properties of ecological communities. Researchers and modelers need to understand the risks of invasion impacts at species, multispecies, and ecosystem levels (Vila et al. 2011). That is the context in which invasions occur.

The challenge to develop scientific methods to monitor the spread, control, and assess the impacts of individual invaders or invasive types (e.g., emergent

PUBLIC DRAFT (7/14/2020)

aquatic vegetation) is ongoing and often boil down to specific situations. Of necessity, most research has focused on correlations such as that between invasive clams and the decline of pelagic fish species. Carefully designed experiments to establish causal relationships are difficult.

There does not seem to be an operational food-web model of the Delta or the necessary components to develop spatially explicit species habitat models through which the impacts of established or potential invaders can be assessed. Assessments of invader impacts are also confounded by other ongoing changes in environmental drivers, so management must be undertaken in the context of a continually changing ecosystem and species pool.

The Dynamic Delta and Non-native Species

How does one manage an ecosystem with an ever-changing species pool? The history of the Delta demonstrates that it is a dynamic place and will become more so in the future. The species pool has undergone continual change. Any new species that becomes established will change the ecosystem in some way. Management must adapt to a continually changing ecosystem and science must be able to forecast future changes and continually evaluate the impacts of a changing species pool on ecosystem structure, function, and services.

The rate, type and impact of new introductions are intertwined with the other major driving forces that change ecosystems, including resource use, climate change, pollution, and habitat alterations (Pysek et al. 2020 and references cited therein). Rapid and accelerating changes in the Delta—the effects of climate change, sea-level rise, changes in water management, salinity intrusion, and so on—will affect virtually all of the factors driving changes in species pools shown in Figure 1. For aquatic invasive species, the changes will affect the vectors and dispersal patterns, characteristics of the receiving habitats, water flows, salinity, seasonal pulses of floods and food-web dynamics, water temperature, and human activities. These will all influence the probability of entry and

PUBLIC DRAFT (7/14/2020)

establishment of invasive species as well as their impacts, creating complex management challenges (Rahel and Olden 2008).

Changing habitats will alter the susceptibility of the Delta to different species and therefore change risk assessment. Pathways may also change. Vulnerability to new nonnatives may differ among habitats and broad taxonomic groups. For example, in a broad meta-analysis, Sorte et al. (2013) found that non-native species were more likely to benefit from the effects of climate change than native species in aquatic ecosystems, but not in terrestrial ones. Non-native fish are generally able to tolerate warmer temperatures, giving them an advantage over native species as the climate warms. Moyle et al. (2013) found that 82% of native fish are vulnerable to the effects of climate change, versus 19% of non-native species.

Consideration of the consequences of predicted climate change in the Delta will be important in forecasting future invasions. Sea-level rise will increase salinity intrusion and inundation in the Delta. Mapping maximum tidal inundation will enable managers to evaluate changes in habitats that will favor the establishment of new species. Climate warming also will change habitat availability. Some species will likely be extirpated from the Delta as their temperature limits are exceeded (e.g. delta smelt), while other species may invade or encroach as higher temperatures or disruptions benefit them, producing the subtractions and additions of species to the species pool shown in Figure 1. Part of this process will involve range expansions of species occurring elsewhere in California.

Warming climate, especially warmer surface water, is expected to shift species distributions and allow non-native species to invade new areas (Walther et al. 2009). Of arguably greater concern are extreme events (e.g., floods, droughts, storms) that will disturb aquatic and wetland ecosystems and facilitate non-native species at every invasion step (Diez et al. 2014). Cloern et al. (2011) modeled how the Delta might change in both average conditions and extreme events.

PUBLIC DRAFT (7/14/2020)

They advised Delta managers to strategize how to adapt to warmer temperatures, higher sea levels, and salinity intrusion and to plan for more runoff in winter and less in spring-summer. They viewed their projections as a starting point, warning “Today's extremes could become tomorrow's norms.”

Changes in temperature and precipitation are expected to affect all aspects of invasion: dispersal pathways (as trade and transport change), establishment (as species ranges shift), impacts (more insect pests, greater food requirements as animals experience stressful conditions, lower stream flows as trees increase evapotranspiration rates), and efforts to manage and control (e.g., shifts in biocontrol-prey interactions, shifts in herbicide tolerance, and more fire-tolerant weeds as drought and fire increase) (Dukes 2011). Along the coast of southern California, invasive non-native plants expand their distributions in years with greater rainfall and lowered soil salinity, which trigger seed germination of upland weedy species as well as native plants (Noe and Zedler 2001a, 2001b; Noe 2002).

Sea levels and climate are expected to change faster than native plants and associated animals can migrate to escape changing conditions. Even a single storm can bring saltwater well inland of normal high tides, killing salt-intolerant species and opening space for invasions. As Callaway and Parker (2012) noted, management of invasive species is already extremely difficult, but “shifting climates will create additional challenges to consider, as changing conditions could create opportunities for a different group of nonnative species, and the future spread of existing invasives will be even more difficult to predict.”

Some invasive species seem pre-adapted to thrive with changing climate. For example, common reed (Box 3, Appendix A) is well adapted to varied climatic conditions where it is native: each lineage has multiple genotypes and grows in diverse habitats and its plastic traits respond to changes related to global warming (temperature, CO₂). Responses to co-occurring environmental changes (drought, salinity, flooding) vary by genotypes within lineages (Skálová et al.

PUBLIC DRAFT (7/14/2020)

2014, Lambertini et al. 2014). As pointed out by Pysek et al. (2020), there are synergies among the interactive drivers affecting new invasions and synergies in the impacts of multiple invaders (e.g., Gaertner et al. 2014).

Recommendations

Changes in the threat assessments for invasive species should be evaluated in the context of a changing environment.

One of the fundamental recommendations from Pysek et al. (2020) is that “Forecasting and scenario development must give more attention to synergies of invasions with climate change and other environmental changes.” We support that recommendation for the Delta. Species distribution data and models and climate models have been used to predict northward movements of fishes under climate change (references). Similar approaches should be tried in the California Delta.

Ecosystem goals must move beyond separate considerations of non-native and native species. Once established, non-native species have become part of the ecosystem.

Climate change scenarios should be incorporated into all management or policy actions regarding non-native species.

Climate warming, sea-level rise, and more extreme environmental conditions will affect all species and habitats in the Delta, accelerating changes in species pools and facilitating the establishment of new non-native species.

A standard climate-change model for the Delta that includes sea-level rise, hydrodynamics, and changes in temperatures should be developed to or enhance the threat assessments for future invaders and changes in populations of current non-native and native species.

Ecological Restoration and Adaptive Management

Findings

- Restoration actions are often accompanied by disturbances that allow invaders to become established.
- Habitat restoration provides the opportunity to use adaptive management approaches to test and select effective methods that favor native species over non-natives.
- Linking non-native species with restoration efforts may enhance the effectiveness of restoration and provide opportunities for adaptive experimentation on control or management approaches.
- Restoration projects can be designed as adaptive-management experiments.

Restoration in the context of Non-native Species

The connection of invasions to restoration is two-fold. First, restorations can create opportunities for non-native species to invade a site, so restorations often include targeted efforts to control or reduce the abundance of non-native species (e.g., by harvesting vegetation). Second, habitat restoration provides the opportunity to use adaptive-management approaches to test and select effective methods that favor native species over non-natives. This includes intentional restoration of invaded sites.

Restoration actions are often accompanied by disturbances that allow invaders to become established. Once non-native plant species become dominant, for example, they often form monotypes that resist eradication. Most attempts to eradicate species covering >1 ha have not achieved their goal (Rejmanek and Pitcairn 2002). The multimillion-dollar attempt to eradicate hybrid cordgrass (*Spartina foliosa* X *S. alterniflora*) along the shores of San Francisco Bay ([San Francisco Estuary Invasive Spartina Project](http://www.spartina.org/): <http://www.spartina.org/>) sparked

PUBLIC DRAFT (7/14/2020)

debates over costs versus. benefits (such as restored habitat for shorebirds, endangered species of rails, or salt marsh harvest mice).

Combinations of co-occurring events and sequences of extreme events may also create opportunities for non-natives to invade restoration sites. Such “sequence events” may have different outcomes when the sequence is reversed (e.g., flood-then-drought effects differ from drought-then-flood effects; Zedler 2010a).

Coinciding extremes, such as the co-occurrence of high river discharge and high coastal water levels, must be considered in risk assessments (Khanna et al. 2018). It is important to include such worst-case scenarios in restoration planning, as there will be surprises and decision protocols will be needed throughout implementation and monitoring.

Restoration often involves transplanting plants into newly restored sites. This may create opportunities for the spread of disease. For example, native plants in northern California nurseries were infected with the non-native fungus, *Phytophthora tentaculata*, which caused root and stem rot. When the native host was planted into restoration sites, the disease spread. While there are now effective guidelines for nurseries to follow, future non-native pathogens await detection (Hunter et al. 2018).

Substantial knowledge is available for replacing non-native plants with former natives. Researchers know where invaders do and do not dominate (Hickson and Keeler-Wolf 2007). Local ecologists often know where there are opportunities to effect control, how to attempt eradication, and what to expect as outcomes. Although preventive programs are envisioned for new invaders, these have not yet been developed or implemented for aquatic invasive plants and wetlands. Inspections, education, and training of people who use Delta waters are essential ingredients of early detection (Ta et al. 2017).

Many researchers with experience in upland vegetation assume that restoring diverse vegetation will resist invasion. Reviews by Guo et al. (2018) and D’Antonio et al. (2016) suggest that aiming for high biodiversity, biomass, and

PUBLIC DRAFT (7/14/2020)

productivity will reduce invasions. However, this is not necessarily true everywhere; Stohlgren et al. (2003) reported the opposite, finding that some diversity hotspots have also been hotspots for invasion.

Restoration projects can be designed as adaptive-management experiments (Zedler 2017). Large adaptive-management experiments can reveal best methods for restoring habitats and managing invaders. Because new invaders will likely appear during restorations, an experimental approach may reveal reasons for their expansions, helping to inform effective management. Adaptive-management experiments may also be the most practical way to determine the effectiveness of new methods to control invasive species, although Conrad et al. (2020) caution that such experimentation may not be possible in some restoration sites because of regulatory restrictions (e.g., protections of endangered species).

In some situations, non-native species may actually benefit ecological restoration. Where non-native species do not unduly threaten other species, ecosystem functioning, or human interests or provide essential ecological or socioeconomic services, they can be tolerated or even used to good advantage (Ewel and Putz 2004). In highly degraded habitats, carefully selected non-native species could be used to accelerate restoration by nitrogen fixation or by acting as nurse plants for native species (Guo et al. 2018). There are always risks where potentially invasive non-native species are involved, but greater risks can be accepted by considering the functional properties of ecosystems rather than using the reconstruction of a biological community as the sole goal of restoration (Ewel and Putz 2004). Both ecosystem functions and the ecology of individual species should be considered in decisions about how (or whether) to manage non-native species.

Ecological restoration is always a long-term process and adaptive management requires monitoring to determine whether and when adaptation of management practices may be necessary. Norton (2009) offered cogent advice: “Restoration

PUBLIC DRAFT (7/14/2020)

outcomes in the face of biological invasions are likely to be novel and will require long-term resource commitment, as any letup in invasive species management will result in the loss of the conservation gains achieved.”

Recommendations

We advocate the formal incorporation of non-native species management and research in ecosystem restoration programs.

Many restoration projects use adaptive management to approach restoration goals as an iterative process. Linking non-native species with restoration efforts may enhance the effectiveness of restoration and provide opportunities for adaptive experimentation on control and management approaches. Proposed restoration efforts should identify pathways for invasive species to enter, implement early detection monitoring, and have an adaptive plan for responding to detections. Setting non-native species goals (like keeping non-native species below 50% of the community) will provide program incentives. When possible, restoration efforts should also take advantage of opportunities to include field experimentation as part of the project design.

Management Coordination and Integration

Our overall recommendation is to **encourage a broader, more forward-looking, integrated approach to non-native species science in the Delta to inform management goals**. Broader means expanding to multiple species and ecosystems; forward-looking means developing predictions and scenarios and forecasting in the context of changing drivers; and integrated means coordinating efforts across interdisciplinary management/enhancement efforts.

Previous recommendations in this review should provide managers with;

- 1) a prioritized list of potential invaders for the immediate and long term that is produced by a robust risk assessment;
- 2) an evaluation of the expected impacts of each high-risk invader;

PUBLIC DRAFT (7/14/2020)

- 3) a monitoring strategy to detect new invaders and map the spread of current invaders; and
- 4) a prioritized list of science actions to help control and understand the impacts of established invaders.

A comprehensive invasive-species coordination plan should be developed for the Delta. The plan should spell out who has the responsibility and how the efforts will be prioritized, supported and funded. Recommendations without responsibilities are unlikely to be implemented (Conrad et al. 2020). Efforts need to be effectively organized and managers are prepared for action. This entails mobilizing the relevant scientific expertise and legal authorities, defining lines of authority, and ensuring that financial and logistical support is sufficient. The wealth of knowledge and experience of Delta managers and researchers is a critical resource that should be brought to bear on future decision making about invasive species in the Delta. The plan should include criteria and performance measures for prioritizing or undertaking control measures by weighing and balancing costs and benefits of non-native or potential invaders and establishing protocols and lines of communication to deal with surprises or the unanticipated arrival of non-natives.

Currently, the Delta Interagency Invasive Species Coordination (DIISC) Team (part of the Sacramento-San Joaquin Delta Conservancy) acts to “foster communication and collaboration among California state agencies, federal agencies, research and conservation groups, and other stakeholders that detect, prevent, and manage invasive species and restore invaded habitats in the Sacramento-San Joaquin Delta”.⁹ DIISC provides a foundation for building broader integration of actions directed toward anticipating, detecting, controlling, and adjusting to invasive species in the Delta. Coordination of monitoring

⁹ [DIISC Team Website](http://deltaconservancy.ca.gov/delta-inter-agency-invasive-species-team/): <http://deltaconservancy.ca.gov/delta-inter-agency-invasive-species-team/>

PUBLIC DRAFT (7/14/2020)

programs, rapid response teams, and management of landscapes and waterscapes to limit invasion corridors cuts across agencies and across species.

An Invasive Species Task Force or Invasive Species Science Center could complement the communication and coordination functions of DIISC by developing a single ‘go to’ science source of expertise and information.

Ultimately, management decisions can be strengthened by using protocols to prioritize actions based, for example, on feasibility, risks, costs, and benefits; by integrating modeling efforts; by testing the effectiveness of new techniques for detecting and controlling invasive species; by developing and using maps of plant and animal biodiversity hotspots and cold spots in the Delta to show where critical functions could be damaged by current or future invasive species; and by incorporating the information and lessons from efforts to deal with invasive species elsewhere and from the growing body of scientific theory and findings about invasive species and their effects.

Conclusions

The species pool of any area within the Delta will continue to change. Some new species will cause measurable harm to the ecology and economy of the Delta. Science can be used to better predict, detect, control, or adapt to these invasive species and inform management to set priorities to minimize harm. Science, however, is only one element among many fiscal, sociological, and political considerations that will ultimately drive allocations of resources to deal with non-native invasive species.

Imagine the following scenario: A particular species (let’s call it “Newtrina”) may be the next invader to the Delta. It enters undetected and become fully established before it is noticed. It disrupts food webs and causes a decline in native species. Management will try to eradicate this species, but it may become permanently established in the Delta and harm ecosystem services valued by people.

PUBLIC DRAFT (7/14/2020)

How should we deal with such a prospect? We should have monitoring systems and food-web studies and spatially explicit habitat models in place to be able to forecast the species' impact and its rate of spread and a central 'go to' base of scientific expertise. We should be able to predict changes in the food web and assess the changes once "Newtrina" has become permanently established. We should be able to tease out the impacts of "Newtrina" relative to ongoing and simultaneous changes in the ecosystem due to climate change, weather extremes, and other driving forces. We should develop protocols for dealing with unanticipated invaders like "Newtrina" that arrive unannounced.

At the present time, the management protocols for preventing, detecting, minimizing the impacts, and adapting to individual invaders are well established and largely adopted at the state and national levels. The science supporting these efforts needs to improve and be applied to the Delta. The approach of focusing on individual invader species has been valuable, although not always effective. However, the rate of invasions and the impact of invaders on ecosystem structure and function are closely linked to other fundamental drivers of ecosystem change, including climate change, resource use, pollution, habitat alteration, and extreme events. Given that the Delta ecosystem has been largely modified, is already highly invaded, and like many other ecosystems is undergoing continual and increasingly rapid change, one might ask: What is the appropriate goal for non-native species management? We can expect that the species pool will continually change and management will need to adapt to the changes. Some of these changes may be predictable and others not. Ultimately management needs to move beyond individual species management to address how to set ecosystem goals in recognition of an ever-changing species pool and uncertainty. The fundamental role of science is to provide management with enough information to set priorities and manage expectations. Developing more forward-looking scientific and predictive science will improve our ability to understand and adapt to those changing conditions.

APPENDIX A: Examples of Significant Non-native Species in the Delta

The following examples highlight several important non-native species and their impacts on the Delta ecosystem. Although non-native species occur throughout the Delta, they have received by far the greatest attention in aquatic environments. The ecological boundaries of upland ecosystems are less well defined relative to the Delta. In agricultural systems, various “pests” and “weeds” (which are also invasive species) have been the focus of intensive prevention and control efforts. While many of our comments apply to non-native and invasive species in any ecosystem, our focus in this report is primarily on aquatic ecosystems.

Bivalves and their effects on the pelagic food web

The Delta has been invaded by several bivalve species that have significantly altered food webs through competition with native filter- and deposit-feeding invertebrates and through altering phytoplankton concentrations. The most notable and well documented of these invaders is *Corbula amurensis*, which was first sighted in the San Francisco Estuary in Grizzly Bay in 1986 (Carlton et al. 1990). The species was likely brought to California as larvae in the ballast of cargo ships. Benthic communities in invaded areas were significantly disrupted and species richness in these habitats gradually decreased during the late 1980s as *C. amurensis* came to dominate the community (Nichols et al. 1990). The combination of the high population growth rate of *C. amurensis* with its filter-feeding efficiency led to a nearly five-fold decrease in average phytoplankton biomass within 2 years of invasion, limiting food availability to zooplankton (Jassby et al. 2002, Thompson 2005). This reduction in phytoplankton biomass shifted food-web dynamics by directing primary production toward benthic consumers (clams) instead of zooplankton (Kimmerer et al. 1994). By depleting native zooplankton, *C. amurensis* facilitated

PUBLIC DRAFT (7/14/2020)

the growth of non-native species in the Delta and shifted the system from a zooplankton community dominated by herbivores and omnivores to one dominated by predatory species. The decreasing food availability for pelagic fish is thought to have contributed to the decline of many fish populations (Nobriga 2002, Cloern and Jassby 2012, Brown et al. 2016). The decrease in productivity of pelagic species stemming from declining phytoplankton was likely due to the combined effects of diversions of freshwater from the Delta, drought conditions that altered salinity and favored non-native zooplankton species, and the *C. amurensis* invasion (Hammock et al. 2019). Thus, the increase in non-native zooplankton in the Delta and associated decline of native pelagic organisms followed multiple human alterations, including water diversions in the Delta (Winder and Jassby 2011, Winder et al. 2011).

Aquatic plants

Several species of non-native aquatic plants reduce native plant diversity and clog waterways, threatening water quality, altering nutrient cycles, and diminishing recreational values in the Delta (Borgnis and Boyer 2016). Of the 19 submerged and floating aquatic plants that occur in the Delta, at least half are non-native. Three of the most widespread non-native species are *Egeria densa* (Brazilian waterweed), *Ludwigia* spp., (water primrose), and *Eichornia crassipes* (water hyacinth) (Khanna et al. 2018a). *Egeria densa* is an example of Submerged Aquatic Vegetation (SAV). It was introduced to the Delta in 1946 from aquarium release and became a species of concern in the 1990s. It forms thick-rooted mats that alter water flow and habitat while impairing recreational activities such as boating and fishing. These hydraulic alterations create a positive feedback loop in which the presence of *E. densa* facilitates its further growth and dispersal (Hestir et al. 2015). The species' low salinity tolerance limits its growth into the western Delta relative to native aquatic vegetation (Borgnis and Boyer 2016). *Egeria densa* cover increased 50% between 2007 and 2014 to about 2900 ha. It is now the dominant submerged aquatic plant, covering 11% of Delta waters (Ustin et al. 2017, Khanna et al. 2015).

PUBLIC DRAFT (7/14/2020)

Eichornia crassipes is an example of a Floating Aquatic Vegetation (FAV) species. It was introduced to California in 1907. It has invaded slow-moving waterways, where its growth changes water quality, displaces native vegetation, clogs channels and marinas, and increases water loss due to its high transpiration rate (Underwood et al. 2006). *Eichornia crassipes* cover increased four-fold between 2004 and 2014 to about 800 ha (Santos et al. 2011a, Dahm et al. 2016). However, 2014 had delayed use of herbicides and it was a peak drought year. Since 2014, water hyacinth cover has been less than it was in 2004-2008 (Ustin et al. 2018).

In addition to *E. densa* and *E. crassipes*, several other non-native plant species pose a threat to Delta waterways. The aquatic alligator weed (*Alternanthera philoxeroides*) is new to the Delta (in 2017, DBW 2017) and is becoming established. It is well known as an aggressive invader in Australia. There, records are available soon after invading for 5 years. During that time, it expanded 4.3 meters per year and produced an average biomass of 4.9 kg dry weight per squared meters per year (Clement et al. 2011). This plant both roots (in shallow water) and produces mats of interwoven of stems that cover waterbodies, restrict human use, exclude native plants, and alter ecosystem functions. *Ludwigia* spp. (water primrose) is a FAV species that increased 4-fold in cover between 2004 and 2016 and encroached into both open water and emergent marsh habitat (Khanna et al. 2018a). *Ludwigia* has been recognized as an emerging problem only in the past decade and now consistently covers more of the waterways than water hyacinth. Coverage in 2014 was similar to that of *E. crassipes* (800 ha) (Boyer and Sutula 2015, Dahm et al. 2016). In 2018 (not considering the south Delta), water primrose occupied about 1200 acres (3.8% of waterways) while water hyacinth was 400 acres (1.3% of waterways) (Ustin et al. 2018). An additional common non-native FAV species of emerging concern, *Limnobium laevigatum* (South American sponge plant), somewhat resembles water hyacinth and is often found alongside it. Common non-native SAV species include *Myriophyllum aquaticum* (parrot feather), *Myriophyllum spicatum* (Eurasian watermilfoil), *Potamogeton crispus* (curlyleaf pondweed), and

PUBLIC DRAFT (7/14/2020)

Cabomba caroliniana (Carolina fanwort) (Ta et al. 2017). *Hydrilla verticillata* (hydrilla) is not yet present in the Delta but occurs elsewhere in California and could migrate into the Delta during high water periods (Ta et al. 2017). Many non-native plant species in the Delta pose major threats to native plant biodiversity, and habitat; species richness of non-native vegetation has been correlated with a decrease in native vegetation species richness and biomass (Santos et al. 2011a).

Despite decades of research and policy directed at managing invasive aquatic plant species, monitoring and controlling their spread remains difficult due to insufficient funding, the absence of consistent monitoring programs, and complex regulations that restrict treatment (Ta et al. 2017). However, monitoring using remote sensing and controls using chemical, mechanical, and biological approaches have been somewhat effective in managing invasive vegetation. For example, several studies have identified and mapped invasive vegetation with high accuracy using hyperspectral remote sensing (Underwood et al. 2006, Hestir et al. 2008, Khanna et al. 2018a). However, this method is subject to error due to spectral variation associated with plant phenology. Nonetheless, remote sensing may be an alternative to costly and time-consuming methods that require direct monitoring of vegetation in remote locations. Drones offer some potential to deliver herbicide to specific patches of invaders ([Huang et al. Project: https://www.ars.usda.gov/research/project/?accnNo=427340](https://www.ars.usda.gov/research/project/?accnNo=427340)).

Efforts to control vegetation may have unintended consequences (Khanna et al. 2012). For example, mechanical shredding of *E. crassipes* may increase overall carbon, nitrogen, and phosphorous levels in the water column up to 10% (Greenfield et al. 2007). Mechanical shredding may also facilitate the spread of many invasive aquatic species, as fragmented plants may re-propagate. Over half of the cut fragments of *E. crassipes* may survive mechanical control and reach a habitat suitable to produce new plants, suggesting that mechanical control may have limited effectiveness in the Delta (Spencer et al. 2006).

PUBLIC DRAFT (7/14/2020)

Alternative uses for the shredded plant material (e.g., feed for livestock) may not be cost effective.

Non-native SAV species also differ functionally from native species. Their greater leaf area, denser canopies, and greater light-use efficiency give them a competitive advantage over native species (Santos et al. 2011b). Thus, the removal of one non-native species may result in colonization by another non-native species instead of the intended native vegetation. Inadvertent effects of control methods must be considered in management of invasive species in the Delta.

Non-native aquatic plants have substantial economic impacts in the Delta, affecting water quality, turbidity (and thus habitat suitability for species such as delta smelt), recreational and commercial boating and fishing, water exports, and virtually all human uses of water. Consequently, there are major ongoing efforts to control invasive plant species in the Delta, spearheaded by a variety of agencies and programs (Box 2). From 2013 to 2017, combined state and federal efforts in chemical control of invasive SAV and FAV averaged approximately \$12.5 million per year (Conrad et al. 2020). Because of regulatory restrictions, control could not be applied everywhere it was needed, and even this level of expenditure was insufficient to achieve effective control of invasive aquatic plants (Conrad et al. 2020).

PUBLIC DRAFT (7/14/2020)

Box 3. Controlling Aquatic Plants

Management of invasive aquatic vegetation in the Delta includes several agencies, including the California Department of Food and Agriculture (CDFA), the California Department of Fish and Wildlife (CDFW), and the California Department of Parks and Recreation, Division of Boating and Waterways (DBW). DBW has the responsibility to control aquatic 'weeds' in the Delta. Because these are independent agencies, coordinating management strategies is often difficult. Several aquatic invasive species, including *E. crassipes* and *E. densa*, are frequently targeted by the DBW Aquatic Invasive Species Program, which is the principal state agency with the authority to treat invasive aquatic species in the Delta (Ta et al. 2017). Treatment typically consists of herbicide application between March and November. Mechanical and biological control measures are also taken to reduce coverage. Biological controls involve alien insects or mites that are introduced to lower the density of non-native vegetation (Ta et al. 2017). Three insect species have been introduced to target *E. crassipes* and two to target *Arundo donax* (giant reed), although only one of these, *Neochetina bruchi* (water hyacinth weevil), has become established in the Delta (Akers et al. 2017, Hopper et al. 2017). There are plans to release other species of weevils and planthoppers in the Delta to selectively feed on invasive vegetation (Ta et al. 2017).

Because managing invasive vegetation is an interagency effort, there are also several collaborative organizations in the Delta that aim to coordinate and manage invasive aquatic species. The Delta Interagency Invasive Species Coordination Team (DIISC) is an interagency group of individuals from agencies focused on preventing, detecting, controlling, and managing invasive species in the Delta (Ta et al. 2017). They aim to increase collaboration among agencies through meeting and facilitating symposia focused on invasive species. USDA sponsors the Delta Region Areawide Aquatic Weed Project (DRAAWP), which focuses on management strategies, control agents, mapping of weeds, and documenting their effects on ecosystem services. DRAAWP centers its efforts on *E. densa*, *E. crassipes*, and *A. donax* and how to best prioritize management practices and provide agencies with essential information.

Wetland vegetation

Wetlands such as tidal or freshwater marshes are a major component of ecological restoration programs in the Delta (e.g. California EcoRestore). Once disturbed, wetlands are vulnerable to invasion by non-native plant species; once established, the invaders are often difficult to control or

PUBLIC DRAFT (7/14/2020)

eradicate. For example, a tall grass called common reed (*Phragmites australis*) is highly invasive in global wetlands and in the Delta, where it crowds out competitors and forms monotypes. Mapping and tracking distributions are difficult in the Delta because native genotypes (not usually invasive) and European strains (highly invasive) both occur and look alike from the air and on the ground (Hickson and Keeler-Wolf 2007). As it does elsewhere, *P. australis* inhabits multiple habitats: palustrine emergent wetlands, freshwater drainage ditches, intertidal bay islands, muted tidal marshes, and wetlands with saline soils (Galatowitsch et al. 1999).

Because wetlands have been a major focus of restoration for a long time, there is considerable knowledge available about several widespread, aggressive invasive plants such as cattails, reed canary grass, and common reed (Zedler and Kercher 2005). Such species reproduce vegetatively from rapidly spreading rhizomes (belowground stems). Their starchy rhizomes serve as reserves that help them resist control using herbicides and cutting and even superficial soil removal. Their tall leaves and stems enable them to outcompete native species. Wetland restoration provides opportunities for field experiments that can enhance our understanding of invader biocontrol methods, herbicide resistance, or the use of heterogeneous topography to facilitate diverse plantings that resist invasions.

Fish

Several studies have substantiated that more non-native than native fish species are present in the Delta and these non-natives have been introduced in a variety of ways.

Many non-native species of fish have been introduced to the Delta through stocking to improve local food and sportfishing opportunities and diversify fish communities. One of the first species introduced was *Alosa sapidissima* (American shad), which was brought to the Sacramento River in 1871 and supported a commercial fishery until the 1950s (Dill and Cordone 1997). *Ameiurus nebulosus* (brown bullhead catfish) were introduced to the San Joaquin

PUBLIC DRAFT (7/14/2020)

River in 1874, followed by several other species of catfish. Striped bass were then introduced to the Carquinez Strait in 1879, leading to a successful commercial fishery that recorded over one million pounds of catch within 20 years. Although large-scale stocking of hatchery-raised striped bass ended in 1992 due to threats to native fish, stocking continued at lower levels in later years.¹⁰

Several other bass species were introduced to California prior to 1900, with records indicating that smallmouth bass (*Micropterus dolomieu*) were first stocked in 1874 and largemouth bass (*Micropterus salmoides*) as early as 1891. Stocking continued for many years. Other bass, including the spotted and redeye bass (*Micropterus punctulatus* and *Micropterus coosae*), were introduced on a lesser scale during the 1930s to 1960s. The establishment of several species of bass in the Delta has resulted in a world-class bass fishery, leading to conflicting goals among individuals managing non-native fish in the Delta: many people wish to recover populations of native species, while others aim to maintain healthy populations of harvestable non-native species. Many of these species, like largemouth and striped bass, prey on or compete with native species like Chinook salmon (*Oncorhynchus tshawytscha*) (Brown and Michniuk 2007). Consequently, management of fish in the Delta involves balancing conflicting interests and ecological goals.

Some other fish species have been introduced as biocontrol agents. *Gambusia affinis* (western mosquitofish) were widely introduced for biological control of mosquitoes in the 1920s. *Menidia audens* (Mississippi silverside) were introduced in the 1960s as a biological control agent; they became widely established by 1975 and are now one of the most widespread and abundant fish species in the Delta (Mahardja et al. 2016).

¹⁰ In February 2020, the California Fish and Game Commission adopted a policy of striving “to maintain a healthy, self-sustaining striped bass population in support of a robust recreational fishery” while eliminating the policy of supporting artificial propagation.

PUBLIC DRAFT (7/14/2020)

Other fish species have been introduced as byproducts of human activity (e.g. see Moyle and Marchetti 2006). One of the most abundant demersal fish in the Delta, *Acanthogobius flavimanus* (yellowfin goby), was first observed in 1963 and was likely introduced through ballast-water transport (Dill and Cordone 1997; Workman and Merz 2007). Their abundance is likely due to their generalist diet, but their inability to reproduce in freshwater has likely limited their expansion. More recent introductions through ballast water include *Tridentiger bifasciatus* (shimofuri goby) and *Tridentiger barbatus* (shokihaze goby), which were first recorded in 1985 and 1997, respectively.

Collectively, non-native species introduced since the 1800s have established populations exceeding the abundance of most native species, resulting in reductions in native fish biodiversity (Moyle 2002; Moyle et al. 2012). In one study that analyzed fish-catch data throughout the Delta between 1994 and 2002, 62% of the species caught and 59% of the overall catch were non-native (Brown and May 2006). Feyrer and Healey (2003) reported that only eight of the 33 species sampled in the southern Delta between 1992 and 1999 were native; no native species accounted for more than 0.5% of the total catch. Higher abundance of native species was correlated with high river flow and turbidity, whereas more non-native fish were associated with warmer water temperatures and low river flow—characteristics of the highly modified south Delta. Similarly, a majority of the overall catch of fish larvae collected between 1990 and 1995 was non-native species associated with low flow and high temperature conditions during the late season; native species were more abundant during early-season conditions (Feyrer 2004). Marchetti et al. (2004) suggested that restoring natural hydrologic processes could mitigate the invasion of non-native fish species while favoring native fish populations.

Historically, the Delta was managed primarily for non-native game fishes, especially striped bass, American shad, and various catishes (Ictaluridae), with some attention also paid to Chinook salmon and steelhead (mainly through hatcheries) and to white sturgeon (Skinner 1962; Kelley 1966; Moyle 2002).

PUBLIC DRAFT (7/14/2020)

Today, formal management of non-native fishes is minimal, even though they contribute substantially to fisheries (e.g. largemouth bass fishery in south and central Delta). Management instead focuses largely on species that are listed under state and federal Endangered Species Acts. However, non-native fishes dominate the fish fauna of the Delta and Suisun Marsh and they form surprisingly integrated fish assemblages with the remaining native species, with a few exceptions (Aguilar-Madrono et al. 2019). This has led Dahm et al. (2019) to suggest that fishes in the Delta should be managed as assemblages with common environmental requirements. For example, striped bass, American shad, delta smelt, and longfin smelt all require a fully functioning estuarine salinity gradient, including substantial outflows to maintain large populations. Historically, all found Suisun Marsh to be an important rearing area.

Non-native fish and submerged aquatic vegetation (SAV)

Both non-native fish and plants have significantly increased in recent decades. Several studies have linked the proliferation of invasive vegetation to the growth of non-native fish populations, but the causal relationship is unclear. One study found that *Egeria densa* is important habitat for juvenile largemouth bass, and the proliferation of this plant likely supported the growth of the largemouth bass fishery in the Delta (Conrad et al. 2016). *Egeria densa* habitat is very productive and several studies have correlated its presence with fish assemblages dominated by non-native species, some of which are predators of native fish such as juvenile salmonids (Brown 2003, Grimaldo et al. 2003, Nobriga et al. 2005, Brown and May 2006, Brown and Michniuk 2007, Loomis 2019). Nobriga et al. (2005) found that native special-status fish species were less abundant in SAV (primarily *E. densa*) habitat than in turbid open water. In contrast, Young et al. (2018) reported that *E. densa* was not correlated with increased macroinvertebrate food for non-native largemouth bass when compared with other SAV species. Although it has been proposed that restoring tidal-wetland habitat would provide important habitat for native fish

PUBLIC DRAFT (7/14/2020)

species, this may only be true where invasive SAV (*E. densa*) is not well established and therefore would not invade the restored habitat (Brown and Michniuk 2007). While restoration for native fish communities looked promising for the northern Delta in 2008, invasive SAV have since increased. For example, Liberty Island was mostly free of SAV in 2008 but now has more than 50% cover of SAV, and the change appears to be persistent (Ustin et al. 2017). Non-native fish might have been facilitated by a concurrent increase in non-native SAV (*Egeria densa*, *Myriophyllum spicatum* and *Potamogeton crispus*). The status and trends of invasive species should be considered when planning future management of both SAV and non-native fish.

Mammals

Nutria are non-native semi-aquatic rodents that are a major threat in the Delta. Although nutria were first introduced to California from South America in 1899 for fur farming, this attempt was commercially unsuccessful (Evans 1970, Carter and Leonard 2002). Subsequent introductions led to a small feral population by the 1940s (Schitoskey 1972), but nutria numbers remained low and the species was eradicated from the state by 1978 (Deems and Pursley 1978). However, a reproducing population was found in the San Joaquin Valley in 2017, and nutria are currently found in the Delta in San Joaquin and neighboring counties (CDFW 2019).

Nutria burrowing and herbivory damage habitats and infrastructure. Nutria burrowing is of great concern in the Delta because levee systems are subject to erosion. Breached levees could allow large agricultural fields to flood, potentially permanently in subsided areas. Nutria feeding is also a threat in the Delta because each animal consumes up to a quarter of its body weight in plants per day. Damage to non-native cattails might not alarm farmers, but they are threatened by losses of rice, corn, and other grains, as well as vegetable crops. Nutria are also vectors for parasites and pathogens. The California multi-agency Response Team is collaborating to eradicate the Delta population. It began as an emergency Incident Command System in 2018 and a formal Nutria Eradication

PUBLIC DRAFT (7/14/2020)

Program in 2019. The Nutria Eradication Program had caught over 1,000 nutria by May 2020 (see Footnote 8).

APPENDIX B: Panelists

To be completed

References

- Aguilar-Medrano, R., Durand, J. R., Cruz-Escalona, V. H., & Moyle, P. B. (2019). Fish functional groups in the San Francisco Estuary: Understanding new fish assemblages in a highly altered estuarine ecosystem. *Estuarine, Coastal and Shelf Science*, 227, 106331.
- Akers, R. P., Bergmann, R. W., & Pitcairn, M. J. (2017). Biological control of water hyacinth in California's Sacramento–San Joaquin River Delta: observations on establishment and spread. *Biocontrol Science and Technology*, 27(6), 755 to 768.
- Anderson, L. W. J. (2005). California's reaction to *Caulerpa taxifolia*: A model for invasive species rapid response. *Biological Invasions* 7, 1003 to 1016.
- Arthur, J. F., Ball, M. D., & Baughman, S. Y. (1996). *Summary of federal and state water project environmental impacts in the San Francisco Bay-Delta Estuary, California*. San Francisco State University.
- Atwood, E. L. (1950). Life history studies of nutria, or coypu, in coastal Louisiana. *Journal of Wildlife Management*, 14(3), 249 to 265.
- Baerwald, M. R., Schreier, B. M., Schumer, G., & May, B. (2012). Detection of the threatened delta smelt in the gut contents of the invasive Mississippi silverside in the San Francisco Estuary using TaqMan assays. *Transactions of the American Fisheries Society*, 141(6), 1600 to 1607.
- Baumsteiger, J., Schroeter, R. E., O'Rear, T. A., Cook, J., & Moyle, P. B. (2017). Long-term surveys show invasive overbite clams (*Potamocorbula amurensis*) are spatially limited in Suisun Marsh, California. *San Francisco Estuary and Watershed Science*, 15(2).
- Beck, K. G., Zimmerman, K., Schardt, J. D., Stone, J., Lukens, R. R., Reichard, S., ... & Thompson, J. P. (2008). Invasive species defined in a policy context: Recommendations from the Federal Invasive Species Advisory Committee. *Invasive Plant Science and Management*, 1(4), 414 to 421.

PUBLIC DRAFT (7/14/2020)

- Borgnis, E., & Boyer, K. E. (2016). Salinity tolerance and competition drive distributions of native and invasive submerged aquatic vegetation in the Upper San Francisco Estuary. *Estuaries and coasts*, 39(3), 707 to 717.
- Boyer, K. E., & Burdick, A. P. (2010). Control of *Lepidium latifolium* (perennial pepperweed) and recovery of native plants in tidal marshes of the San Francisco Estuary. *Wetlands ecology and management*, 18(6), 731 to 743.
- Boyer, K., & Sutula, M. (2015). Factors controlling submersed and floating macrophytes in the Sacramento-San Joaquin Delta. Southern California Coastal Water Research Project. Technical Report 870. Costa Mesa, CA.
- Boyer, K., Miller, J., Borgnis, E., Patten, M., & Moderan, J. (2016). Salinity effects on native and introduced submerged aquatic vegetation of Suisun Bay and the Delta. Final Report to the CALFED Ecosystem Restoration Program and California Department of Fish and Wildlife. Romberg Tiburon Center for Environmental Studies, San Francisco State University Grant No.: E1183007/ERP-11-S11.
- Brown, L. R. (2003). Will tidal wetland restoration enhance populations of native fishes?. *San Francisco Estuary and Watershed Science*, 1(1).
- Brown, L. R., Kimmerer, W., Conrad, J. L., Lesmeister, S., & Mueller-Solger, A. (2016). Food webs of the Delta, Suisun Bay, and Suisun Marsh: An update on current understanding and possibilities for management. *San Francisco Estuary and Watershed Science*, 14(3).
- Brown, L. R., & May, J. T. (2006). Variation in spring nearshore resident fish species composition and life histories in the lower San Joaquin watershed and delta. *San Francisco Estuary and Watershed Science*, 4(2).
- Brown, L. R., & Michniuk, D. (2007). Littoral fish assemblages of the alien-dominated Sacramento-San Joaquin Delta, California, 1980–1983 and 2001–2003. *Estuaries and Coasts*, 30(1), 186 to 200.
- California State Parks, Division of Boating and Waterways (DBW). (2017). Floating aquatic vegetation control program 2017 annual monitoring report.
- Callaway, J. C., & Parker, V. T. (2012). Current issues in tidal marsh restoration. Chapter 18 (pp. 252-262). *Ecology, Conservation, and Restoration of Tidal Marshes*; University of California Press: Berkeley, CA, USA, 233 to 252.
- Carlton, J. T. (1996). Biological invasion and cryptogenic species. *Ecology*, 77(6), 1653.
- Carlton, J. T., Thompson, J. K., Schemel, L. E., & Nichols, F. H. (1990). Remarkable invasion of San Francisco Bay (California, USA) by the Asian clam

PUBLIC DRAFT (7/14/2020)

Potamocorbula amurensis. I. Introduction and dispersal. *Marine Ecology Progress Series*, 66, 81 to 94.

Carter, J. & Leonard, B.P. (2002). Review of the literature on the worldwide distribution, spread of, and efforts to eradicate the coypu (*Myocastor coypus*): *Wildlife Society Bulletin*, 30(1), 162 to 175.

Charles, S. P., Kominoski, J. S., Troxler, T. G., Gaiser, E. E., Servais, S., Wilson, B. J., ... & Kelly, S. (2019). Experimental saltwater intrusion drives rapid soil elevation and carbon loss in freshwater and brackish Everglades marshes. *Estuaries and Coasts*, 42(7), 1868 to 1881.

Clements, D., Dugdale, T. M., & Hunt, T. D. (2011). Growth of aquatic alligator weed (*Alternanthera philoxeroides*) over 5 years in south-east Australia. *Aquatic Invasions*, 6(1), 77 to 82.

Cloern, J. E., & Jassby, A. D. (2012). Drivers of change in estuarine-coastal ecosystems: Discoveries from four decades of study in San Francisco Bay. *Reviews of Geophysics*, 50(4).

Cloern, J. E., Knowles, N., Brown, L. R., Cayan, D., Dettinger, M. D., Morgan, T. L., ... & Jassby, A. D. (2011). Projected evolution of California's San Francisco Bay-Delta-River system in a century of climate change. *PloS one*, 6(9), Article e24465.

Cohen, A. N., & Carlton, J. T. (1995). Nonindigenous aquatic species in a United States estuary: A case study of the biological invasions of the San Francisco Bay and Delta. Connecticut Seagrant, Groton, CT.

Cohen, A. N., & Carlton, J. T. (1998). Accelerating invasion rate in a highly invaded estuary. *Science*, 279(5350), 555 to 558.

Colautti, R. I. & MacIsaac, H. J. (2004). A neutral terminology to define "invasive" species. *Diversity and Distributions*, 10, 134 to 141.

Conrad, J. L., Bibian, A. J., Weinersmith, K. L., De Carion, D., Young, M. J., Crain, P., ... & Sih, A. (2016). Novel species interactions in a highly modified estuary: Association of Largemouth Bass with Brazilian waterweed *Egeria densa*. *Transactions of the American Fisheries Society*, 145(2), 249 to 263.

Conrad, J. L., Chapple, D., Bush, E., Hard, E., Caudill, J., Madsen, J. D., ... & Khanna, S. (2020). Critical needs for control of invasive aquatic vegetation in the Sacramento-San Joaquin Delta. Report to the Delta Stewardship Council, May 2020.

Crooks, J. A. (2002). Characterizing ecosystem-level consequences of biological invasions: The role of ecosystem engineers. *Oikos*, 97(2), 153 to 166.

PUBLIC DRAFT (7/14/2020)

D'Antonio, C.M., August-Schmidt, E., & Fernandez-Going, B. (2016). Invasive species and restoration challenges. In M.A. Palmer, J.B. Zedler, & D.A. Falk (Eds.), *Foundations of Restoration Ecology* (pp. 216-244). Washington DC: Island Press.

Dahm, C. N., Parker, A. E., Adelson, A. E., Christman, M. A., & Bergamaschi, B. A. (2016). Nutrient dynamics of the Delta: Effects on primary producers. *San Francisco Estuary and Watershed Science*, 14(4).

Dahm, C., Kimmerer, W., Korman, J., Moyle, P. B., Ruggerone, G. T., & Simenstad, C. A. (2019). Developing Biological Goals for the Bay-Delta Plan: Concepts and Ideas from an Independent Scientific Advisory Panel. A final report to the Delta Science Program. Sacramento: Delta Stewardship Council.

Davis, M. A. (2011). Don't judge species on their origins. *Nature*, 474, 153 to 154.

Deems, E. F., & Pursley, D. (1978). North American furbearers: Their management, research and harvest status in 1976: International Association of Fish and Wildlife Agencies in cooperation with the Maryland Department of Natural Resources-Wildlife Administration: College Park, University of Maryland.

Dick, J. T., Laverty, C., Lennon, J. J., Barrios-O'Neill, D., Mensink, P. J., Britton, J. R., ... & Dunn, A. M. (2017). Invader Relative Impact Potential: A new metric to understand and predict the ecological impacts of existing, emerging and future invasive alien species. *Journal of Applied Ecology*, 54(4), 1259 to 1267.

Diez et al. 2014

Dill, W. A., & Cordone, A. J. (1997). History and status of introduced fishes in California, 1871-1996. *Fish Bulletin*, 178, 1 to 414.

Dukes 2011

Durand, J., Fleenor, W., McElreath, R., Santos, M. J., & Moyle, P. B. (2016). Physical controls on the distribution of the submersed aquatic weed *Egeria densa* in the Sacramento-San Joaquin Delta and implications for habitat restoration. *San Francisco Estuary and Watershed Science*. 14(1).

Eller, F., Skálová, H., Caplan, J. S., Bhattarai, G. P., Burger, M. K., Cronin, J. T., ... & Lambertini, C. (2017). Cosmopolitan species as models for ecophysiological responses to global change: The common reed *Phragmites australis*. *Frontiers in plant science*, 8, 1833.

Eller, F., Lambertini, C., Nguyen, L. X., & Brix, H. (2014). Increased invasive potential of non-native *Phragmites australis*: Elevated CO₂ and temperature alleviate salinity effects on photosynthesis and growth. *Global change biology*, 20(2), 531 to 543.

PUBLIC DRAFT (7/14/2020)

Elton, C. S. (1958). *The ecology of invasions by animals and plants*. University of Chicago Press.

Evans, J. (1970). *About nutria and their control* (No. 86). US Fish and Wildlife Service.

Ewel, J.J., & Putz, F.E. (2004). A place for alien species in ecosystem restoration. *Frontiers in Ecology and the Environment*, 2(7), 354 to 360.

Feyrer, F. (2004). Ecological segregation of native and alien larval fish assemblages in the southern Sacramento-San Joaquin Delta. In *American Fisheries Society Symposium* (pp. 67 to 80). American Fisheries Society.

Feyrer, F., & Healey, M. P. (2003). Fish community structure and environmental correlates in the highly altered southern Sacramento-San Joaquin Delta. *Environmental Biology of Fishes*, 66(2), 123 to 132.

Fleenor, W.E., Hanak, E., Lund, J.R. & Mount J. (2008). Delta hydrodynamics and water salinity with future conditions. Appendix C. Comparing Futures for the Sacramento-San Joaquin Delta. Public Policy Institute of California, David, CA.

Foxcroft, L. C. (2009). Developing thresholds of potential concern for invasive alien species: hypotheses and concepts. *Koedoe*, 51(1).

Gaertner, M., Biggs, R., Te Beest, M., Hui, C., Molofsky, J., & Richardson, D. M. (2014). Invasive plants as drivers of regime shifts: Identifying high-priority invaders that alter feedback relationships. *Diversity and Distributions*, 20, 733 to 744.

Galatowitsch, S. M., Anderson, N. O., & Ascher, P. D. (1999). Invasiveness in wetland plants in temperate North America. *Wetlands*, 19(4), 733 to 755.

Greenfield, B. K., Siemering, G. S., Andrews, J. C., Rajan, M., Andrews, Jr., S. P., & Spencer, D. F. (2007). Mechanical shredding of water hyacinth (*Eichornia crassipes*): Effects on water quality in the Sacramento-San Joaquin River Delta, California. *Estuaries and Coasts*, 30(4), 627 to 640.

Grimaldo, L. F., Miller, R. E., Peregrin, C. M., & Hymanson, Z. P. (2003). Spatial and temporal distribution of native and alien ichthyoplankton in three habitat types of the Sacramento-San Joaquin Delta. In *American Fisheries Society Symposium* (pp. 81 to 96). American Fisheries Society.

Godoy, O. (2019). Coexistence theory as a tool to understand biological invasions in species interaction networks: Implications for the study of novel ecosystems. *Functional Ecology*, 33, 1190 to 1201.

PUBLIC DRAFT (7/14/2020)

Guo, Q., Brockway, D. G., Larson, D. L., Wang, D., & Ren, H. (2018). Improving ecological restoration to curb biotic invasion - a practical guide. *Invasive Plant Science and Management*, 11(4), 163 to 174.

Hammock, B. G., Moose, S. P., Solis, S. S., Goharian, E., & Teh, S. J. (2019). Hydrodynamic modeling coupled with long-term field data provide evidence for suppression of phytoplankton by invasive clams and freshwater exports in the San Francisco Estuary. *Environmental management*, 63(6), 703 to 717.

Hestir, E. L., Khanna, S., Andrew, M. E., Santos, M. J., Viers, J. H., Greenberg, J. A., Rajapakse, S. S., & Ustin, S. L. (2008). Identification of invasive vegetation using hyperspectral remote sensing in the California Delta ecosystem. *Remote Sensing of Environment*, 112(11), 4034 to 4047.

Hestir et al. 2015

Hickson, D., Keeler-Wolf, T., & Region, B. D. (2007). Vegetation and land use classification and map of the Sacramento-San Joaquin River Delta. Report prepared for Bay Delta Region of California Department of Fish and Game. *Vegetation Classification and Mapping Program, California Department of Fish and Game, Sacramento, CA.*

Holl, K. D., Howard, E. A., Brown, T. M., Chan, R. G., de Silva, T. S., Mann, E. T., ... & Spangler, W. H. (2014). Efficacy of exotic control strategies for restoring coastal prairie grasses. *Invasive Plant Science and Management*, 7(4), 590 to 598.

Hopper, J. V., Pratt, P. D., McCue, K. F., Pitcairn, M. J., Moran, P. J., & Madsen, J. D. (2017). Spatial and temporal variation of biological control agents associated with *Eichhornia crassipes* in the Sacramento-San Joaquin River Delta, California. *Biological Control*, 111, 13 to 22.

Hosler, D. M. (2017). Where is the body? Dreissenid mussels, raw water testing, and the real value of environmental DNA. *Management of Biological Invasions*, 8, 335 to 341.

Hui, C. A., Rudnick, D., & Williams, E. (2005). Mercury burdens in Chinese mitten crabs (*Eriocheir sinensis*) in three tributaries of southern San Francisco Bay, California, USA. *Environmental Pollution*, 133, 481 to 487.

Hunter, S., Williams, N., McDougal, R., Scott, P., & Garbelotto, M. (2018). Evidence for rapid adaptive evolution of tolerance to chemical treatments in *Phytophthora* species and its practical implications. *PloS one*, 13(12), e0208961.

Invasive Species Symposium 2019 White Paper.

PUBLIC DRAFT (7/14/2020)

IPBES (Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services). (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services. Bonn, Germany:

Janovsky R.M., & Larson E.R. (2019). Does invasive species research use more militaristic language than other ecology and conservation biology literature? *NeoBiota* 44, 27 to 38.

ecology and conservation biology literature? *NeoBiota* 44: 27–38. Jassby, A. D., Cloern, J. E., & Cole, B. E. (2002). Annual primary production: Patterns and mechanisms of change in a nutrient-rich tidal ecosystem. *Limnology and Oceanography*, 47(3), 698 to 712.

Jerde, C., Mahon, A. R., Chadderton, W. L., & Lodge, D. M. (2011). “Sight-unseen” detection of rare aquatic species using environmental DNA. *Conservation Letters*, 4, 150 to 157.

Khanna, S., Santos, M. J., Hestir, E. L., & Ustin, S. L. (2012). Plant community dynamics relative to the changing distribution of a highly invasive species, *Eichhornia crassipes*: a remote sensing perspective. *Biological Invasions*, 14(3), 717 to 733.

Khanna, S., Bellvert, J., Shapiro, K., & Ustin, S. L. (2015). Invasions in State of the Estuary 2015: Status and Trends Updates on 33 Indicators of Ecosystem Health. Retrieved from Oakland, California, USA.

Khanna, S., Santos, M. J., Boyer, J. D., Shapiro, K. D., Bellvert, J., & Ustin, S. L. (2018a). Water primrose invasion changes successional pathways in an estuarine ecosystem. *Ecosphere*, 9(9), 1 to 18.

Khanna S, Conrad JL, Caudill J, Christman M, Darin G, Ellis D, Gilbert P, Hartman R, Kayfetz K, Pratt W, et al. (2018b). Framework for Aquatic Vegetation Monitoring in the Delta. Aquatic Vegetation Project Work Team, Interagency Ecological Program.

Khanna et al. 2019

Kimmerer, W. J., Gartside, E., & Orsi, J. J. (1994). Predation by an introduced clam as the likely cause of substantial declines in zooplankton of San Francisco Bay. *Marine Ecology Progress Series*, 113, 81 to 93.

Kolar, C. S. & Lodge, D. M. (2001). Progress in invasion biology: predicting invaders. *Trends in Ecology and Evolution*, 16(4), 199 to 204.

Leidy, R. A., & Fiedler, P. L. (1985). Human disturbance and patterns of fish species diversity in the San Francisco Bay drainage, California. *Biological Conservation*, 33(3), 247 to 267.

PUBLIC DRAFT (7/14/2020)

- Liao, C, Peng, R., Luo, Y., Zhou, X., Xiaowen, W., Fang, C, ... & Li, B. (2007). Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytologist*, 177(3), 706 to 714.
- Loomis, C. M. (2019). Density and distribution of piscivorous fishes in the Sacramento–San Joaquin Delta. Thesis.
- Lund, J.R., Hanak, E., Fleenor, W.E., Bennett, W.A., Howitt, R.E., Mount, J.F., and Moyle, P.B. (2010). Comparing Futures for the Sacramento San Joaquin Delta. University of California Press, Berkeley, CA.
- Luoma, S. N., Dahm, C. N., Healey, M., & Moore, J. N. (2015). Challenges facing the Sacramento–San Joaquin Delta: Complex, chaotic, or simply cantankerous?. *San Francisco Estuary and Watershed Science*, 13(3).
- Mahardja, B., Conrad, J. L., Lusher, L., & Schreier, B. (2016). Abundance trends, distribution, and habitat associations of the invasive Mississippi silverside (*Menidia audens*) in the Sacramento–San Joaquin Delta, California, USA. *San Francisco Estuary and Watershed Science*, 14(1).
- Mann, C.C. (2011). *Uncovering the New World Columbus Created*. New York, NY, A.A. Knopf.
- MacNally, R., Thomson, J.R., Kimmerer, W.J., Feyrer, F., Newman, K.B., Sih, A., Bennett, W.A., Brown, L., Fleishman, E., Culberson, S.D., & Castillo, G. (2010). Analysis of pelagic species decline in the upper San Francisco Estuary using multivariate autoregressive modeling (MAR). *Ecological Applications*, 20(5), 1417 to 1430.
- Marchetti, M. P., Light, T., Moyle, P. B., & Viers, J. H. (2004). Fish invasions in California watersheds: Testing hypotheses using landscape patterns. *Ecological Applications*, 14(5), 1507 to 1525.
- Marchetti, M. P., Moyle, P. B., & Levine, R. (2004). Alien fishes in California watersheds: Characteristics of successful and failed invaders. *Ecological Applications*, 14(2), 587 to 596.
- Matern, S. A., & Brown, L. R. (2005). Invaders eating invaders: Exploitation of novel alien prey by the alien shimofuri goby in the San Francisco Estuary, California. *Biological Invasions*, 7, 497 to 507.
- Mihaylov, R., Dimitrov, R., Binev, R., & Stamatova-Yovcheva, K. (2017). A study of some biological, anatomical and related environmental features of nutria /*Myocastor coypus*/ from the territory of Stara Zagora region. *Veterinary Journal of Mehmet Akif Ersoy University*, 2(1), 7 to 15.
- Millennium Ecosystem Assessment (2005).

PUBLIC DRAFT (7/14/2020)

Moyle, P. B. (2011). [Striped bass control: Cure worse than disease?](https://californiawaterblog.com/2011/01/31/stripped-bass-control-the-cure-worse-than-the-disease/) [Blog post]. Retrieved January 31, 2011 from <https://californiawaterblog.com/2011/01/31/stripped-bass-control-the-cure-worse-than-the-disease/>

Moyle, P. B. (2020). [Striped Bass: An important indicator species in the Delta.](https://californiawaterblog.com/2020/01/12/stripped-bass-an-important-indicator-species-in-the-delta/) [Blog post]. Retrieved January 12, 2020 from <https://californiawaterblog.com/2020/01/12/stripped-bass-an-important-indicator-species-in-the-delta/>

Moyle, P.B., & Bennett, W.A. (2008). The future of the Delta ecosystem and its fish. Appendix D. Comparing Futures for the Sacramento-San Joaquin Delta. Public Policy Institute of California, Davis, CA

Moyle, P. B., Hobbs, J., & O'Rear, T. (2012). Fishes. In A. Palaima (Ed.), Ecology, conservation and restoration of tidal marshes: The San Francisco Estuary (1st ed., pp. 161 to 173). Berkeley: University of California Press.

Moyle, P. B., & Marchetti, M. P. (2006). Predicting invasion success: Freshwater fishes in California as a model. *Bioscience*, 56(6), 515 to 524.

Nichols, F. H., Thompson, J. K., & Scheme, L. E. (1990). *Potamocorbula amurensis*. 11. Displacement of a former community. *Marine Ecology Progress Series*, 66, 95 to 101.

Nobriga, M. L. (2002). Larval delta smelt diet composition and feeding incidence: environmental and ontogenetic influences. *California Fish and Game*, 88(4), 149 to 164.

Nobriga, M. L., Feyrer, F., Baxter, R. D., & Chotkowski, M. (2005). Fish community ecology in an altered river delta: Spatial patterns in species composition, life history strategies, and biomass. *Estuaries*, 28(5), 776 to 785.

Noe, G. B. (2002). Temporal variability matters: Effects of constant vs. varying moisture and salinity on germination. *Ecological Monographs*, 72(3), 427 to 443.

Noe, G. B., & Zedler, J. B. (2001a). Spatio-temporal variation of salt marsh seedling establishment in relation to the abiotic and biotic environment. *Journal of Vegetation Science*, 12(1), 61 to 74.

Noe, G. B., & Zedler, J. B. (2001b). Variable rainfall limits the germination of upper intertidal marsh plants in southern California. *Estuaries*, 24(1), 30 to 40.

Norton, D. A. (2009). Species invasions and the limits to restoration: Learning from the New Zealand experience. *Science*, 325(5940), 569 to 571.

PUBLIC DRAFT (7/14/2020)

Perino, A., Pereira, H. M., Navarro, L. M., Fernández, N., Bullock, J. M., Ceaușu, S., ... & Pe'er, G. (2019). Rewilding complex ecosystems. *Science*, 364(6438), Article eaav5570.

Pyšek, P., & Richardson, D. M. (2010). Invasive species, environmental change and management, and health. *Annual review of environment and resources*, 35, 25 to 55.

Pyšek, P., Hulme, P.E., Simberloff, D., Bacher, S., Blackburn, T.M, Carlton, J.T., Dawson, W., & Richardson, D.M. (2020). Scientists' warming on invasive alien species. *Biological Reviews*.

Rahel, F.J., & Olden, J. D. (2008). Assessing the effects of climate change on aquatic invasive species. *Conservation Biology*, 22, 521 to 533.

Rejmánek, M., & Pitcairn, M. J. (2002). When is eradication of exotic pest plants a realistic goal? In C.R. Veitch and M.N. Clout (Eds.), *Turning the Tide: The Eradication of Invasive Species* (pp. 249–253). Gland, Switzerland and Cambridge, UK International Union for Conservation of Nature and Natural Resources.

Ricciardi, A. & Rasmussen, J. B. (1998). Predicting the identify and impact of future biological invaders: a priority for aquatic resource management. *Canadian Journal of Fisheries and Aquatic Sciences*, 55, 1759 to 1765.

Ricciardi, A., & Simberloff, D. (2009). Assisted colonization is not a viable conservation strategy. *Trends in ecology & evolution*, 24(5), 248 to 253.

Roy, K., Jablonski, D., Valentine, J. W. (2002). Body size and invasion success in marine bivalves. *Ecology Letters*, 5(2), 163 to 167.

Rudnick, D., Halat, K., & Resh, V. H. (2000). Distribution, ecology and potential impacts of the Chinese mitten crab (*Eriocheir sinensis*) in San Francisco Bay. University of California Water Resources Center Contribution 206. (pp. 74).

Rudnick, D. & Resh, V. H. (2002). A survey to examine the effects of the Chinese mitten crab on commercial fisheries in Northern California. *IEP Newsletter*, 15, 19 to 21.

Rudnick, D., Hieb, K., Grimmer, K., & Resh, V. H. (2003). Patterns and processes of biological invasion: the Chinese mitten crab in San Francisco Bay. *Basic and Applied Ecology*, 4, 249 to 262.

Rudnick, D., Chan, V. & Resh, V. H. (2005). Morphology and impacts of the burrows of the Chinese mitten crab, *Eriocheir sinensis* H. Milne Edwards (Decapoda, Grapsidae) in south San Francisco Bay, California, U.S.A. *Crustaceana*, 78(7), 787 to 807.

PUBLIC DRAFT (7/14/2020)

- Ruiz, G. M., Fofonoff, P. W., Carlton, J. T., Wonham, M. J., & Hines, A. H. (2000). Invasion of coastal marine communities in North America: Apparent patterns, processes, and biases. *Annual review of ecology and systematics*, 31(1), 481 to 531.
- Ruiz, G. M., Fofonoff, P. W., Steves, B., Foss, S. F., & Shiba, S. N. (2011). Marine invasion history and vector analysis of California: A hotspot for western North America. *Diversity and Distributions*, 17(2), 362 to 373.
- Santos, M. J., Anderson, L. W., & Ustin, S. L. (2011a). Effects of invasive species on plant communities: An example using submersed aquatic plants at the regional scale. *Biological Invasions*, 13(2), 443 to 457.
- Santos, M. J., Hestir, E. L., Khanna, S., & Ustin, S. L. (2011b). Image spectroscopy and stable isotopes elucidate functional dissimilarity between native and nonnative plant species in the aquatic environment. *New Phytologist*, 193(3), 683 to 695.
- Scianni, C., Celballos Osuna, L., Dobroski, N., Falkner, M., Thompson, J., & Nedelcheva, R. (2019). 2019 Biennial report on the California marine invasive species program. Report produced for the California state legislature.
- SFEI-ASC (San Francisco Estuary Institute-Aquatic Science Center. (2014). A Delta Transformed: Ecological Functions, Spatial Metrics, and Landscape Change in the Sacramento-San Joaquin Delta. Prepared for the California Department of Fish and Wildlife and Ecosystem Restoration Program. A Report of SFEI-ASC's Resilient Landscape Program, Publication #729, San Francisco Estuary Institute-Aquatic Science Center, Richmond, CA.
- Shrader-Frechette, K. (2001). Non-indigenous species and ecological explanation. *Biology and Philosophy*, 16, 507 to 519.
- Simberloff, D. & Rejmanek, M. (Eds.). (2011). *Encyclopedia of Biological Invasions*. Berkeley, CA, University of California Press.
- Sorte, C. J. B., Ibáñez, I., Blumenthal, D. M., Molinari, N. A., Miller, L. P., Grosholz, E. D.,... & Dukes, J. S. (2013). Poised to prosper? A cross - system comparison of climate change effects on native and non - native species performance. *Ecology Letters*, 16, 261 to 270.
- Sousa, R., Gutiérrez, J. L., & Aldridge, D. C. (2009). Non-indigenous invasive bivalves as ecosystem engineers. *Biological Invasions*, 11(10), 2367 to 2385.
- Spencer, D. F., Ksander, G. G., Donovan, M. J., Liow, P. S., Chan, W. K., Greenfield, B. K., ... & Andrews, S. P. (2006). Evaluation of water hyacinth survival and growth in the Sacramento Delta, California, following cutting. *Journal of aquatic plant management*, 44(1), 50 to 60.

PUBLIC DRAFT (7/14/2020)

Srèbalienė, G., Olenin, S., Minchin, D., & Narščius, A. (2019). A comparison of impact and risk assessment methods based on the IMO Guidelines and EU invasive alien species risk assessment frameworks. *PeerJ*, 7, Article e6965.

Stoeckle, B. (2016). Environmental DNA as a monitoring tool for the endangered freshwater pearl mussel (*Margaritifera margaritifera* L.): a substitute for classical monitoring approaches?. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(6), 1120 to 1129.

Stohlgren, T. J., Barnett, D. T., & Kartesz, J. C. (2003). The rich get richer: Patterns of plant invasions in the United States. *Frontiers in Ecology and the Environment*, 1(1), 11 to 14.

Ta, J., Anderson, L. W. J., Christman, M. A., Khanna, S., Kratville, D....& Viers, J. H. (2017). Invasive aquatic vegetation management in the Sacramento-San Joaquin River Delta: Status and recommendations. *San Francisco Estuary and Watershed Science*, 15(4), 1 to 19.

Talley, T. S., Crooks, J. A., & Levin, L. A. (2001). Habitat utilization and alteration by the invasive burrowing isopod, *Sphaeroma quoyanum*, in California salt marshes. *Marine Biology*, 138(3), 561 to 573.

Thompson, J. K. (2005). One estuary, one invasion, two responses: Phytoplankton and benthic community dynamics determine the effect of an estuarine invasive suspension-feeder. In *The comparative roles of suspension-feeders in ecosystems* (pp. 291 to 316). Springer, Dordrecht.

Ustin, S. L., Khanna, S., Lay, M., Shapiro, K., & Ghajarnia, N. (2017). Enhancement of Delta Smelt (*Hypomesus transpacificus*) habitat through adaptive management of invasive aquatic weeds in the Sacramento-San Joaquin Delta. Report submitted to California Department of Water Resources.

Ustin, S. L., Khanna, S., Lay, M., & Shapiro, K. (2018). Enhancement of Delta Smelt (*Hypomesus transpacificus*) habitat through adaptive management of invasive aquatic weeds in the Sacramento-San Joaquin Delta. Report submitted to California Department of Water Resources.

Underwood, E. C., Mulitsch, M. J., Greenberg, J. A., Whiting, M. L., Ustin, S. L., & Kefauver, S. C. (2006). Mapping invasive aquatic vegetation in the Sacramento-San Joaquin Delta using hyperspectral imagery. *Environmental Monitoring and Assessment*, 121(1 to 3), 47 to 64.

Vilà, M., Espinar, J. L., Hejda, M., Hulme, P. E., Jarošík, V., Maron, J. L., ... & Pyšek, P. (2011). Ecological impacts of invasive alien plants: A meta-analysis of their effects on species, communities and ecosystems. *Ecology letters*, 14(7), 702 to 708.

PUBLIC DRAFT (7/14/2020)

Walther, G. R., Roques, A., Hulme, P. E., Sykes, M. T., Pyšek, P., Kühn, I., ... & Czucz, B. (2009). Alien species in a warmer world: risks and opportunities. *Trends in Ecology & Evolution*, 24(12), 686 to 693.

Whipple, A.A., Grossinger, R.M., Rankin, D., Stanford, B., & Askevold, R.A. (2012). Sacramento-San Joaquin Delta Historical Ecology Investigation: Exploring Pattern and Process. Prepared for the California Department of Fish and Wildlife and Ecosystem Restoration Program. A Report of SFEI-ASC's Historical Ecology Program, SFEI-ASC Publication #672, San Francisco Estuary Institute-Aquatic Science Center, Richmond, CA.

Wiens, J. D., Anthony, R. G., & Forsman, E. D. (2014). Competitive interactions and resource partitioning between northern spotted owls and barred owls in western Oregon. *Wildlife Monographs*, 185, 1 to 50.

Willner, G. R., Chapman, J. A., Pursley, D. (1979). Reproduction, physiological responses, food habits, and abundance of nutria on Maryland marshes. *Wildlife Monographs*, 65, 3 to 53.

Winder, M., & Jassby, A. D. (2011). Shifts in zooplankton community structure: Implications for food web processes in the upper San Francisco Estuary. *Estuaries and Coasts*, 34(4), 675 to 690.

Winder, M., Jassby, A. D., & Mac Nally, R. (2011). Synergies between climate anomalies and hydrological modifications facilitate estuarine biotic invasions. *Ecology letters*, 14(8), 749 to 757.

Wolff, W. J. (1998). Exotic invaders of the meso-oligohaline zone of estuaries in the Netherlands: Why are there so many?. *Helgoländer Meeresuntersuchungen*, 52(3), 393.

Woo, I., & Takekawa, J. Y. (2012). Will inundation and salinity levels associated with projected sea level rise reduce the survival, growth, and reproductive capacity of *Sarcocornia pacifica* (pickleweed)?. *Aquatic botany*, 102, 8 to 14.

Workman, M. L., & Merz, J. E. (2007). Introduced yellowfin goby, *Acanthogobius flavimanus*: Diet and habitat use in the lower Mokelumne River, California. *San Francisco Estuary and Watershed Science*, 5(1).

Young, M. J., Conrad, J. L., Bibian, A. J., & Sih, A. (2018). The effect of submersed aquatic vegetation on invertebrates important in diets of juvenile largemouth bass *Micropterus salmoides*. *San Francisco Estuary and Watershed Science*, 16(2).

Zedler, J.B. (2010). Tussock sedge: A restoration superplant? *Arboretum Leaflet* 22.

PUBLIC DRAFT (7/14/2020)

Zedler, J. B. (2017). What's new in adaptive management and restoration of coasts and estuaries? *Estuaries and Coasts*, 40(1), 1 to 21.

Zedler, J. B., & Kercher, S. (2005). Wetland resources: Status, trends, ecosystem services, and restorability. *Annual Review of Environment and Resources*, 30, 39 to 74.

Zedler, J. B., & Stevens, M. L. (2018). Western and traditional ecological knowledge in ecocultural restoration. *San Francisco Estuary and Watershed Science*, 16(3), 1 to 18.